

## REVIEW

## Epigenetics and locust life phase transitions

Ulrich R. Ernst, Matthias B. Van Hiel, Geert Depuydt, Bart Boerjan, Arnold De Loof and Liliane Schoofs\*

## ABSTRACT

Insects are one of the most successful classes on Earth, reflected in an enormous species richness and diversity. Arguably, this success is partly due to the high degree to which polyphenism, where one genotype gives rise to more than one phenotype, is exploited by many of its species. In social insects, for instance, larval diet influences the development into distinct castes; and locust polyphenism has tricked researchers for years into believing that the drastically different solitary and gregarious phases might be different species. Solitary locusts behave much as common grasshoppers. However, they are notorious for forming vast, devastating swarms upon crowding. These gregarious animals are shorter lived, less fecund and transmit their phase characteristics to their offspring. The behavioural gregarisation occurs within hours, yet the full display of gregarious characters takes several generations, as does the reversal to the solitary phase. Hormones, neuropeptides and neurotransmitters influence some of the phase traits; however, none of the suggested mechanisms can account for all the observed differences, notably imprinting effects on longevity and fecundity. This is why, more recently, epigenetics has caught the interest of the polyphenism field. Accumulating evidence points towards a role for epigenetic regulation in locust phase polyphenism. This is corroborated in the economically important locust species *Locusta migratoria* and *Schistocerca gregaria*. Here, we review the key elements involved in phase transition in locusts and possible epigenetic regulation. We discuss the relative role of DNA methylation, histone modification and small RNA molecules, and suggest future research directions.

**KEY WORDS:** Locust phase, Polyphenism, Locust swarming, *Locusta migratoria*, *Schistocerca gregaria*, *Apis mellifera*, Invertebrate, DNA methylation, Histone modification, Methyloome

## Introduction

The term epigenetics tends to take a variety of meanings (Haig, 2004; Jablonka and Lamb, 2002). In its narrow sense, it can be defined as ‘meiotically and mitotically heritable changes in gene expression, not based on DNA sequence alterations’ (Riggs et al., 1996). In a broader sense, it can also be interpreted as ‘modifications of chromosome structure’ (Bird, 2007). Independent of the interpretation, however, epigenetics did not receive much attention in insect research until recent years.

The most prominent epigenetic mechanisms, namely (1) methylation of cytosine in DNA, (2) modifications of histone proteins and (3) nucleosome positioning and regulation by non-coding RNA, were only rarely the focus of the entomology field. This may have been due to the general low, often almost undetectable, levels of methylation in insects and other invertebrates

(Glastad et al., 2011), including the prime model organisms *Drosophila melanogaster* and *Caenorhabditis elegans*, suggesting that DNA methylation in insects is of minor importance (Glastad et al., 2011). However, it should be noted that the detection of methylated bases in DNA and/or RNA is prone to errors, especially when the methylation rate is relatively low. As a result, methylation levels, or even its occurrence per se, have been heavily debated in *C. elegans* (Klass et al., 1983; Simpson et al., 1986) and *D. melanogaster* (Achwal et al., 1983; Lyko et al., 2000; Raddatz et al., 2013).

The discovery of a functional DNA methylation system in the European honeybee *Apis mellifera* (Wang et al., 2006) triggered a renewed interest in the role of epigenetics in insect biology. In addition, recent advances in sequencing technologies have tremendously facilitated the systematic study of epigenomes and epigenetics in a wide range of insects (Beeler et al., 2014; Bonasio et al., 2012; Krauss et al., 2009; Lo et al., 2012; Lyko et al., 2010; Smith et al., 2012; Walsh et al., 2010; Weiner et al., 2013; Xiang et al., 2013; Ye et al., 2013; Zemach et al., 2010; Zwieter et al., 2012).

The diversity of epigenetic systems in insects has made them interesting models for understanding DNA methylation (Lyko and Maleszka, 2011). The observation that locust DNA may be the subject of slightly higher methylation levels (1.3–1.9% of total cytosines, depending on the tissue) than reported for other insect species (Boerjan et al., 2011) revived interest in epigenetic control in locust phase polyphenism. Polyphenism, where one genotype produces several phenotypes, is relatively common in the Animal Kingdom (Simpson et al., 2011). Popular examples in vertebrates include temperature-dependent sex determination in some fish and reptiles, where the ambient temperature experienced during a specific time in development triggers the development into male or female (Navarro-Martí et al., 2011). The insect clade provides some of the very best models for polyphenism (reviewed in Moczek, 2010; Simpson et al., 2011; Whitman and Ananthakrishnan, 2009). Metamorphosis in holometabolic insects, where larva, pupa and imago often differ dramatically in various traits, is just one example. Other well-studied examples include the formation of eye-spots in the butterfly *Bicyclus anynana* (reviewed in Brakefield and Frankino, 2009), or the striking difference in horn size in dung beetles (*Onthophagus*) (Kijimoto et al., 2013). In aphids it has been shown that both wing polyphenism – i.e. the development of winged or wingless morphs – and morph polyphenism – i.e. oviparity or viviparity – are maternally regulated (Hartfelder and Emlen, 2012; Ogawa and Miura, 2014; Zera and Denno, 1997). In addition, research into polyphenism and developmental plasticity has often relied on the caste phenomenon in social insects.

Locusts undergo similar drastic changes when they start swarming, a behaviour that has been extensively studied for many years (Burrows et al., 2011; Pener and Simpson, 2009; Wang and Kang, 2014). When solitary locusts become gregarious, they form enormous groups of countless individuals, spanning occasionally hundreds of square kilometres (Ferenz, 1990). The socio-economic impact of these swarms is estimated to be up to several billion US

Functional Genomics and Proteomics Lab, KU Leuven, Naamsestraat 59, bus 2465, B-3000 Leuven, Belgium.

\*Author for correspondence (Liliane.Schoofs@bio.kuleuven.be)

### List of abbreviations

AKH	adipokinetic hormone
APRP	adipokinetic hormone precursor-related peptide
Cas	CRISPR-associated protein
CNS	central nervous system
CpG	cytosine followed by guanine
CRISPR	clustered regularly interspaced short palindromic repeat
Dnmt	DNA methyltransferase
EST	expressed sequence tag
HAT	histone acetyl transferase
HDAC	histone deacetylase
HDM	histone demethylase
HMT	histone methyltransferase
HSP	heat-shock protein
JH	juvenile hormone
JHPH	juvenile hormone binding protein, hexamerins, prophenoloxidase and haemocyanins
LITE	light-inducible transcriptional effector
MBD	methyl-binding protein
miRNA	microRNA
ncRNA	non-coding RNA
NPP	neuroparsin precursors
piRNA	PIWI-interacting RNA
PTM	post-translational modifications
RNAi	RNA interference
RRBS	reduced representation bisulphite sequencing
siRNA	short interfering RNA
TALE	transcription activator-like effector
TALLEN	transcription activator-like effector nuclease
ZFN	zinc finger nuclease

dollars ([www.fao.org/docrep/018/i2940e/i2940e17.pdf](http://www.fao.org/docrep/018/i2940e/i2940e17.pdf)). In this review, we will focus on the epigenetic aspects of polyphenetic transitions in the most important and best-studied species, the migratory locust *Locusta migratoria* and the desert locust *Schistocerca gregaria*.

### Comparing solitary and gregarious phases: morphology, behaviour and physiology

Phase transition between the solitary and the gregarious form encompasses extreme phenotypic plasticity at multiple levels including locust morphology, behaviour, neurochemistry and physiology (Pener and Simpson, 2009; Pener and Yerushalmi, 1998; Uvarov, 1966; Verlinden et al., 2009) (Fig. 1; supplementary material Table S1). Among the most obvious effects of phase transition are changes in morphological appearance including body size and colour (Pener and Simpson, 2009; Uvarov, 1966). Solitary individuals are generally bigger and cryptically coloured compared with their long-term gregarious counterparts, which have a conspicuous bright body colour (Fig. 2). More subtle anatomical differences can be seen in the shape and size of the eyes, wings, antennae and jumping hindlegs, as well as in the distribution of sensory receptors (Pener, 1991; Pener and Yerushalmi, 1998). In addition, phase transition induces a broad range of physiological differences in lifespan, metabolism, immune responses, endocrinology and reproductive physiology (Pener and Yerushalmi, 1998; Verlinden et al., 2009; Wang and Kang, 2014). Higher fecundity and smaller eggs have been observed in solitary versus gregarious forms of desert and migratory locusts (Maeno and Tanaka, 2008; Maeno and Tanaka, 2009). Increased population density also alters the rate of sexual maturation, but the effects are species dependent: gregarious desert locusts sexually mature more rapidly, whereas the opposite has been reported for *L. migratoria* (Maeno and Tanaka, 2009; Norris, 1952; Norris, 1950). In gregarious male desert locusts, this is accompanied by a bright




**Fig. 1. Photomontage of a solitary (left) and gregarious (right) last instar *Schistocerca gregaria*.** Crowding induces gregarisation and is instigated via (1) mechanical stimulation of the hindlegs, and/or (2) the combined sight and smell of other locusts. Gregarisation results in altered morphology, physiology, behaviour and colour (only the last of these is depicted; the right half of the image has been modified to artificially match the colours of gregarious locusts). Original image courtesy of Tom Fayle, modified by Fabian Ernst.

yellow coloration due to the incorporation of yellow protein into the cuticle (Sas et al., 2007; Wybrandt and Andersen, 2001).

Behavioural dissimilarities are the most intriguing phase-related differences. Solitary locusts normally avoid each other, but increased population density can rapidly trigger attraction to other locusts, resulting in aggregation behaviour that can lead to the generation of devastating migratory swarms (Pener and Simpson, 2009; Rogers et al., 2003; Uvarov, 1966). Gregarious morphs exhibit a wider dietary range, display increased locomotory activity, and will fly predominantly during the daytime, in contrast to isolated locusts, which generally fly at night (Pener, 1991; Uvarov, 1977).

Related to behavioural changes imposed by group living and foraging, phase transition in locusts has been found to alter brain structure and functioning. Long-term gregarious desert locusts have a smaller body size, but their brain is substantially larger – about 30% – than that of long-term solitary locusts (Ott and Rogers, 2010). In addition, Ott and Rogers reported that the relative distribution of brain regions differs between the two phases. Solitary locusts invest more in lower level sensory processing, reflected by their relatively large primary olfactory and visual neuropils. In contrast, the larger brains of gregarious locusts are more dedicated to the integration of sensory cues in higher level processing regions, which is thought to support their lifestyle as generalist foragers in dense, migratory swarms where competition among group members is high (Ott and Rogers, 2010). Other changes in brain functioning situate at the level of circuit activity and function (Ayali et al., 2004; Blackburn et al., 2010; Burrows et al., 2011; Fuchs et al., 2003). For example, gregarious *L. migratoria* show reduced habituation of interneuron activity in response to approaching objects. Phase transition also affects associative learning in the desert locust (Simões et al., 2013). Long-term solitary locusts learn more quickly to associate an odour with an aversive food source, an effect that can be overridden by crowding, enabling locusts to adopt a different feeding strategy. Furthermore, brain functioning is affected by changes in the neurochemistry of the locust's nervous system upon



**Fig. 2. Body coloration of *S. gregaria* depends largely on phase and developmental stage.** Solitary locusts are larger and cryptically coloured; gregarious locusts display aposematic colours. Males are shown in the gregarious phase as last instar larva (A) and imago (B), and in the solitary phase as last instar larva (C) and imago (D).

phenotypic transformation (Anstey et al., 2009; Rogers et al., 2004; Verlinden et al., 2009).


Solitary locusts can switch to gregarious behaviour in only a few hours (Ellis, 1953; Ellis, 1962), whereas other changes, in colour, morphology and reproductive physiology, alter on a much slower time scale (Pener and Simpson, 2009; Pener and Yerushalmi, 1998; Roessingh et al., 1993; Simpson et al., 1999). Reversible transition between the two phases occurs gradually and through intermediate forms, spanning multiple generations. Rapid behavioural changes can rely on short-term neuronal plasticity to alter circuit activity and function, whereas morphological responses such as in brain structure or muscle morphology depend on long-term remodelling that can span several generations (Burrows et al., 2011; Simpson and Miller, 2007; Tanaka and Maeno, 2006). This suggests that epigenetic signals accumulate when reinforced, or fade away with time when they are not reinstalled, which would account for the slow change of some phase characters (Burggren, 2015; Jablonka and Raz, 2009) (Fig. 3).

### Making the switch: initiation of phase transition

To date, no single factor has been identified that can induce the entire set of alterations in a locust during phase transition. However, during the past two decades, a number of factors have been identified that are involved.

### Visual, olfactory and/or mechanosensory information

Two distinct sensory pathways are involved in the onset and continuation of aggregating behaviour: the cerebral pathway, prompted by the combined visual and olfactory stimuli, and the thoracic pathway, induced by tactile information (Burrows et al., 2011). Stimulating the hindleg, but not other parts of the body, is sufficient for gregarisation behaviour to occur (Ellis, 1959; Simpson et al., 2001). In the Australian plague locust, *Chortoicetes terminifera*, stimulation of the antennae, but not the hindlegs, is sufficient to elicit a phase transition (Cullen et al., 2010). Not only tactile stimulation but also the combined sight and smell of conspecific or heterospecific locusts can induce gregariousness in



**Fig. 3. Hypothetical model for epigenetic remodelling in locust phase transitions.** Crowding causes profound differences in neuronal and hormonal signalling, gene expression and epigenetic modifications that eventually lead to significant changes in behaviour, physiology and morphology on different time scales. Hormones, gene expression and epigenetic modifications influence each other. Epigenetic marks will perpetuate these changes over moultings and egg formation. Eggs may be primed in the ovary and in the egg pod by an egg foam factor. When offspring also experience crowding, epigenetic alterations may accumulate that subsequently lead to morphological changes and the phenotype of long-term gregarious locusts.



locust nymphs as well (Leo Lester et al., 2005; Roessingh et al., 1998). However, it has been reported that certain contact chemicals by themselves can induce gregarisation behaviour (Heifetz et al., 1997; Heifetz et al., 1998; Heifetz et al., 1996). For a review on the roles of semiochemicals in locusts, we refer the reader to Hassanali and co-workers (Hassanali et al., 2005). In contrast, Tanaka and Nishide report that the sight of moving animals is sufficient to induce darkening in desert locusts, whereas odour had barely any effect (Tanaka and Nishide, 2012).

Tactile information through the antennae reflecting the degree of crowding experienced by the mother directly influences the colour of hatchlings in *S. gregaria* and *L. migratoria*, arguing for a maternal factor (Maeno et al., 2011). It is likely that an alkylated L-DOPA analogue recently isolated from egg foam, which was found to induce gregarious behaviour in nymphs hatched from treated eggs deposited by solitary females, fulfils this role (Islam, 2013; Miller et al., 2008).

### Neuroendocrine control of phase transition: corazonin, juvenile hormone, serotonin and dopamine

Although the concentration of several potential neurochemicals differs between solitary and gregarious locusts, only serotonin shows a dramatic transient increase within hours of crowding (Rogers et al., 2004). Moreover, injection of serotonin (and its analogues) is sufficient to induce gregariousness in both *S. gregaria* and *L. migratoria*, whereas injecting serotonin antagonists inhibits tactile stimulation-induced phase transition (Anstey et al., 2009; Ma et al., 2011). These observations establish serotonin as an important regulatory agent in phase transition, an effect that is probably mediated through protein kinase A signalling (Ott et al., 2012). We note that other researchers report no influence on attraction/avoidance behaviour by serotonin (Tanaka and Nishide, 2013) or on darkening of hatchlings (Maeno et al., 2011) in *S. gregaria*. However, in the latter experiments, serotonin was injected through the abdominal sternites in the thorax instead of directly into the thoracic ganglia, which could account for the difference in results.

Surprisingly, injection of serotonin or serotonin receptor (5-HT<sub>2</sub>) agonists into the head cavity of gregarious *L. migratoria* nymphs shifts their behaviour towards solitariousness (Guo et al., 2013a), suggesting serotonin is involved in both the gregarisation and solitarisation decision. These results correspond well with earlier observations that serotonin displays a peak in the optic lobes (located in the head cavity) shortly after isolation, while a similar peak is seen in the thoracic ganglion when these individuals are re-aggregated (Rogers et al., 2004).

Pharmacological and transcriptional silencing experiments identified dopamine, in addition to serotonin, as another potent gregarisation factor in the migratory locust, driving both behavioural transition and melanin deposition (Ma et al., 2011).

Tanaka has postulated that juvenile hormone (JH) in conjunction with corazonin can account for body colour polyphenism, in both the desert and migratory locust (Tanaka, 2006). Corazonin, an undecapeptide released from the corpora cardiaca, causes darkening of locust body colour (Vandersmissen et al., 2006), whereas JH and JH analogues induce the green body colour in solitary forms (Hasegawa and Tanaka, 1994). Besides instigating dark body colouring, corazonin induces a more convex pronotum in isolated-reared locusts (Tanaka et al., 2002), changes body aspect ratios towards those of gregarious animals (Breuer et al., 2003; Hoste et al., 2002; Maeno et al., 2004; Tanaka et al., 2002) and reduces the number of antennal sensilla to that of crowd-reared locusts (Maeno

and Tanaka, 2004; Yamamoto-Kihara et al., 2004). However, corazonin cannot induce gregarious behaviour (Hoste et al., 2003) or the darkening of offspring of solitary individuals (Tanaka, 2001).

### Comparing solitary and gregarious phases: molecular differences derived from -omics data

The recent publication of the first draft of the 6.5 Gbp genome sequence of the migratory locust (Wang et al., 2014) marks a crucial milestone that should greatly aid researchers in their quest to unravel the molecular foundations underlying locust phase change. Earlier -omics approaches have been instrumental in gaining new insights into the physiological transformation that differentiates the solitary locust from its gregarious phase, and will be summarised in this section.

#### Proteomics and peptidomics

*Schistocerca gregaria* pacifastin-like precursors (SGPP1–4), peptides with still unknown functions, display a strong phase-dependent transcription in the brain and fat body, suggesting a physiological role in phase transition (Simonet et al., 2005; Simonet et al., 2004). Similarly, *L. migratoria* LMPP2 has a differential expression between the two phases (Kang et al., 2004). Neuroparsin precursors (NPP) from the desert locust (*Scg-NPPI–4*) similarly show a very distinct phase-dependent expression (Claeys et al., 2006b; Claeys et al., 2005). Interestingly, NPP transcription is induced upon injection with JH or 20-hydroxyecdysone (Claeys et al., 2006a).

In *L. migratoria*, higher levels of neuroparsin A (NP-A) and ovary maturing parsin (Lom-OMP) were found in the corpora cardiaca of crowded locusts. In contrast, the concentration of adipokinetic hormone (AKH) precursor-related peptide (APRP) was decreased (Ayali et al., 1996). APRPs are by-products of AKH synthesis with an as-yet unknown functional role (Baggerman et al., 2002; Hatle and Spring, 1999). Neuropeptide profiles of the corpora cardiaca and haemolymph confirmed phase-differentiating levels of AKH I and II and APRPs in *S. gregaria*, but strong sex- and age-dependent concentration differences were also observed (Clynen et al., 2002).

The most enigmatic factor is the phase-related peptide (Clynen et al., 2002), a 6 kDa peptide present in much higher concentrations in the haemolymph of gregarious locusts (up to 0.1 mmol l<sup>-1</sup>) compared with solitary locusts. Interestingly, upon isolation of crowd-reared locusts, this peptide shows a progressive decrease in concentration spanning multiple generations. This finding represents another clear manifestation of the epigenetic aspects of locust phase differentiation and maintenance. Higher concentrations of this peptide are also found in the eggs of gregarious than in the eggs of solitary *S. gregaria* (Rahman et al., 2003a; Rahman et al., 2002).

#### Metabolomics

Comparison of the metabolic profiles of haemolymph between solitary and crowd-reared *S. gregaria* using <sup>1</sup>H NMR spectroscopy revealed over 20 differentially abundant metabolites (Lenz et al., 2001). Most notably, crowd-reared insects showed much lower levels of lipid and trehalose, as well as the polyamine putrescine, a breakdown product of amino acids.

More recently, HPLC-GC/MS-based metabolic profiles of solitary and gregarious *L. migratoria* haemolymph were analysed over the time course of phase transition (Wu et al., 2012). Over 45% (319) of the detected metabolites differed between the two phases, with multiple lipids, carbohydrates, amino acids and carnitines showing

the most prominent changes. Interestingly, the genetic or pharmacological manipulation of acylcarnitine and acetylcarnitine levels induces both metabolic and behavioural changes associated with phase transition. As carnitine is responsible for the transport of fatty acids into the mitochondrial matrix prior to their degradation, this could reflect increased energy requirements in gregarious locusts. Alterations in energy metabolism, membrane fluidity and lipid-mediated cell signalling are all potential avenues by which changes in lipid metabolism could be implicated in the phase transition.

### Transcriptomics

Rahman and co-workers used differential display RT-PCR (Rahman et al., 2003b) in an early attempt to investigate differential expression in phase transition. They found one gene with higher expression in gregarious desert locusts (resembling *Drosophila* SPARC) and one unidentified gene with higher expression in solitary locusts. The subsequent construction of expressed sequence tag (EST) databases has greatly facilitated locust research. In 2004, a high-coverage EST data set (76,012 ESTs) of the whole body, head, hindlegs and midgut tissue of *L. migratoria* hoppers was generated (Kang et al., 2004). A general repression of (mostly anabolic, biosynthetic and muscle-specific) gene expression was observed in the hindleg, midgut and head tissue of gregarious compared with solitary animals, an observation consistent with their weaker leaping ability. The relative expression pattern in the head of gregarious animals was characterised by the strong upregulation of a functionally diverse set of genes. Most notably, very strong activation of the JHPH (juvenile hormone binding protein, hexamerins, prophenoloxidase and haemocyanins) gene family was observed in the head of gregarious locusts. These findings suggest extensive expressional changes in nerve cells that could reflect how hormonal signals govern the phase state transition. The same group also described elevated transcript levels of six heat-shock protein (HSP) genes in gregarious locusts compared with solitary locusts (Wang et al., 2007). This activation of HSP expression probably reflects a stress response triggered by increased population density. This increase in the expression of genes related to stress is consistent with the findings of Badisco and co-workers; they detected 214 differentially expressed genes (Badisco et al., 2011b) in an EST database (34,672 ESTs) generated from the central nervous system (CNS) of both phases of the desert locust (Badisco et al., 2011a). The upregulation of HSP genes and immunity genes in gregarious locusts seems to offer protection from the acute detrimental effects of crowding. At the same time, a reduction in the expression of genes associated with energy metabolism and protein synthesis occurs. This reduction of metabolism and biosynthesis in gregarious locusts was also found in migratory locusts by comparing the transcriptome (72,977 sequences) of developing solitary and gregarious locusts (Chen et al., 2010). Taken together, these data clearly suggest that gregarious locusts are associated with a physiological stress state (Boerjan et al., 2010). Conversely, expression of antioxidant genes is repressed in gregarious desert locusts, suggesting increased susceptibility to oxidative stress (Badisco et al., 2011b). In gregarious migratory locusts, a cluster of synaptic transport components, neurotransmitter and neuromodulator receptors, neurotransmitter synthetases and G protein-coupled receptors (GPCRs) was also prominently upregulated, pointing to an important neuromodulatory aspect in phase transition (Chen et al., 2010). In addition, it was found that fourth instar locusts displayed the most divergence between polymorphic states. This was confirmed by micro-array studies,

which illustrated that the top pathways affected in gregarious nymphs at fourth stage were involved in general metabolism and catecholamine biosynthesis (Ma et al., 2011).

Another micro-array study on fourth instar migratory locusts during phase transition revealed differential expression in chemosensory proteins and *takeout* proteins (Guo et al., 2011). Genes of these families were found to be highly expressed in peripheral tissues (sensilla, antennae, labial palps, wings and hindlegs) but not internal tissues. RNA interference (RNAi) against chemosensory proteins increased repulsion in gregarious fourth instar nymphs, whereas knockdown of one *takeout* gene (*LmigT01*) increased attraction behaviour in solitary fourth instar nymphs, probably by altering peripheral olfactory sensitivity.

Large-scale transcriptome analysis revealed 105 retro-elements in the migratory locust, some of which show a differential expression pattern between the solitary and gregarious phase at the fifth instar and in adults (Jiang et al., 2012). The developmental and tissue-specific expression pattern of a single class I transposon element was also highly different in gregarious compared with solitary locusts (Guo et al., 2010). These observations have prompted the Kang lab to propose a regulatory role for these transposon elements in the phase transition of migratory locusts (Jiang et al., 2012). Interestingly, the degree of genome methylation constitutes an important factor determining the transcriptional activity of (retro)transposons and suggests one possible role for genome methylation in the modulation of phenotypic plasticity in insects (Slotkin and Martienssen, 2007).

A recent transcriptome analysis of brain tissue of locusts experiencing short-term (64 h) gregarisation and solitarisation revealed a staggering 4893 differentially expressed genes (28% of the transcriptome) in the two processes (Wang et al., 2014). Increased expression of genes in synaptic transmission, carbohydrate metabolism and nucleosome assembly seems to point towards increased neuronal activity during locust crowding, while at the same time the lowered expression of genes in redox biology suggests suppression of antioxidative responses in the CNS. The same authors also identified 45 genes where alternative splicing results in differential isoform expression between phases. A common theme in the transcriptome, alternative splicing and methylome datasets (see below) is the differential expression (and methylation) of cytoskeletal/microtubular genes, probably reflecting the neuronal plasticity accompanying the behavioural changes upon phase transition.

### Epigenetic mechanisms in invertebrates

#### Histones in insect epigenetics

Next to DNA methylation and demethylation processes, reversible post-translational modifications (PTMs) of histone proteins are the best-studied elements of epigenetic mechanisms. More than 160 histone modifications, e.g. methylation, acetylation and phosphorylation (Bannister and Kouzarides, 2011; Suganuma and Workman, 2011; Tan et al., 2011), alter chromatin structure and density and hence the accessibility of DNA, which influences transcription rates ('histone code') (Jenuwein and Allis, 2001; Strahl and Allis, 2000). Histone modifications and chromatin states have been intensively studied in *D. melanogaster* (Filion et al., 2010; Kharchenko et al., 2011; Nègre et al., 2011). Less is known about the role of histone PTMs in other insects. However, Nanty and colleagues showed that patterns of histone PTMs are largely conserved between invertebrate species and can therefore be predicted for different taxa (Nanty et al., 2011). Indeed, DNA methylation and histone modifications seem to work together, if not

redundantly, to influence gene expression patterns (Hunt et al., 2013a; Hunt et al., 2013b).

Several histone PTMs have been characterised in honeybees (Dickman et al., 2013). To date, the best evidence for the involvement of histone modifications in insect polyphenism comes from the carpenter ant *Camponotus floridanus*, where the pattern of acetylation of lysine 27 at histone 3 (H3K27ac) differs between males, and major and minor workers (Simola et al., 2013).

#### Non-coding RNA and other epigenetic mechanisms

While DNA methylation and histone modifications are the most prominent epigenetic mechanisms studied to date, non-coding (nc)RNA and heritable protein alteration have recently received increased attention. Altered protein conformations transmittable to subsequent generations have been studied in yeasts (Halfmann et al., 2010; Halfmann et al., 2012; Halfmann and Lindquist, 2010). Similar mechanisms have been suggested for *Drosophila*, but have yet to be proven (Tariq et al., 2013).

ncRNAs may have a more prominent role in epigenetic mechanisms than previously thought. New classes of ncRNA, including short interfering RNA (siRNA), microRNA (miRNA), PIWI-interacting RNA (piRNA) and long ncRNA, have varying roles in gene regulation (Jacquier, 2009; Moazed, 2009; Pauli et al., 2011). In honeybees and ants, for instance, miRNAs have been implicated in (temporal) caste differences (Behura and Whitfield, 2010; Bonasio et al., 2010; Greenberg et al., 2012; Guo et al., 2013b; Liu et al., 2012).

#### Epigenetics in life phase transitions

Epigenetic mechanisms could play a crucial role in life phase transitions in insects. In the honeybee, DNA methylation and histone modifications mark two important processes: (1) the irreversible differentiation of a female larva into a queen or worker phenotype (Kucharski et al., 2008) and (2) the reversible shift for worker bees from a temporal nurse subcaste to the forager subcaste (Herb et al., 2012; Lockett et al., 2012). This has also been studied in several ant and wasp species (Bonasio, 2014; Bonasio et al., 2012; Simola et al., 2013; Weiner et al., 2013) [see also Bonasio (Bonasio, 2015) in the current issue]. The differentiation into a queen or a worker has dramatic consequences: a honeybee queen lives several years, is much larger, highly fertile and also differs in many more morphological traits and behavioural characteristics from her sisters that developed into workers and have a life expectancy of only a few weeks (Winston, 1987).

Despite the overwhelming indications for an important role of epigenetics in the regulation of phase transitions in insects, the direct evidence is relatively limited. The best-known example is the induction of queen-like phenotypes in honeybees, *Apis mellifera*, by downregulating of DNA methyltransferase 3 (Dnmt3) (Kucharski et al., 2008; Li-Byarlay et al., 2013). In the buff-tailed bumblebee, *Bombus terrestris*, experimental alteration of DNA methylation by feeding with 5-aza-20-deoxycytidine (decitabine) renders queenless worker bees more aggressive and more fertile (Amarasinghe et al., 2014). However, the drug treatment was only successful in callow workers (i.e. younger than 1 day), whereas older workers are also able to activate their ovaries.

In the crustacean water flea *Daphnia magna*, exposure to 5-azacytidine reduced overall DNA methylation as well as body length (Vandegheuchte et al., 2010). Interestingly, this hypomethylation pattern was transferred to two subsequent generations that were not exposed to the drug, demonstrating transgenerational epigenetic inheritance. These two generations were also shorter, but as yet there

is no link established between hypomethylation and body length. However, it should be noted that 5-azacytidine and other nucleoside analogues are not specific DNA methylation inhibitors and also affect other pathways (Gnyszka et al., 2013; Poirier et al., 2014).

Further evidence for epigenetic inheritance in *Drosophila* and *Daphnia* is reviewed elsewhere (Youngson and Whitelaw, 2008), but the mechanisms of this 'soft' inheritance are generally not well understood.

#### Evidence for epigenetics in locusts

##### DNA methylation

To our knowledge, the earliest study on DNA methylation in locusts dates from 1951 (Wyatt, 1951), and reported that 0.96% of all cytosines are methylated in *L. migratoria*. Surprisingly, these early findings were followed by a 60 year gap. In 2011, we showed that, compared with that of other insects, *S. gregaria* DNA is relatively highly methylated (1.3–1.9% of total cytosines, depending on the tissue) (Boerjan et al., 2011). The *Schistocerca* transcriptome contains transcripts for some of the enzymes belonging to the epigenetic machinery, including a methyl binding protein (MBD), a histone acetyl transferase (HAT), a histone deacetylase (HDAC) and homologues of Dnmt1 and Dnmt2 (Boerjan et al., 2011; Falckenhayn et al., 2013). Expression levels of Dnmt2 in the metathoracic ganglion change during crowding. Next-generation shotgun bisulphite sequencing to identify the *S. gregaria* methylome confirmed 1.3–1.4% cytosine methylation, 90% of which are in a CpG (cytosine followed by guanine) context (Falckenhayn et al., 2013). The locust genome is more highly methylated than most known insect genomes [but less so than the genomes of other Orthoptera: *L. migratoria*, 1.6% (Wang et al., 2014); *Grylloptarpa fossor*, 3% (Sarkar et al., 1992); *Chorthippus parallelus*, 4.06±0.68% (Lechner et al., 2013)]. As the genome of *S. gregaria* has not been sequenced so far, the sequences were mapped against an EST database (Badisco et al., 2011a). Of those that could be mapped, 3.2% and 3.1% of cytosines for brain and metathoracic ganglia, respectively, are methylated, 97% of which are in a CpG context. This more than twofold higher methylation level in comparison with the overall methylation pattern suggests that methylation is targeted to exons. In contrast to the honeybee (Lyko et al., 2010) and the silkworm (Xiang et al., 2010; Zemach et al., 2010), but similar to the stick insect (Krauss et al., 2009) and *L. migratoria* (Robinson et al., 2011; Wang et al., 2014), repetitive elements (ribosomal DNA, transposons) are also methylated. Genes with a low CpG observed/expected ratio are assumed to have been (historically) methylated in the germline, as methylated cytosines tend to mutate to thymines, causing depletion of cytosines over evolutionary time scales (Bird, 1980; Duncan and Miller, 1980). Indeed, these genes are more methylated in *L. migratoria*: 20% of the contigs were over 95% methylated and another 20% were more than 65% methylated, an unusually high methylation rate and distinct from other invertebrates. However, gene methylation did not correlate with gene expression levels in six of the investigated genes, suggesting that there is no straightforward link between methylation levels and gene expression.

Recently, the early report of DNA methylation in *L. migratoria* (Wyatt, 1951) was confirmed by Robinson and co-workers, who found that 1.3% of the total cytosines are methylated (Robinson et al., 2011), which is in the same range as *S. gregaria*. Similar to *S. gregaria*, DNA methylation in *L. migratoria* is not restricted to gene bodies, but also targeted to repetitive elements (Robinson et al., 2011). In addition, Robinson and colleagues found transcripts for MBD 2/3, Dnmt2 and two copies of Dnmt1 (Robinson et al., 2011).



Interestingly, genes differentially expressed between the two phases in *L. migratoria* show signs of CpG depletion. The hypermutability of methylated cytosines leads to the formation of thymines via deamination. Thus, a depletion of CpGs over time occurs if highly methylated sequences in the germline were affected. The CpG O/E value is the ratio between observed and expected CpGs within a sequence and is a signature of historical DNA methylation in the germline.

In 2014 the genome sequence of *L. migratoria* was reported (Wang et al., 2014) and confirmed the findings of Robinson and co-workers (Robinson et al., 2011). The *L. migratoria* genome encodes an apparently functional methylation system, containing two copies of Dnmt1 and a single copy of Dnmt2 and 3. Besides the genes reported by Wang and co-workers, we found evidence of additional genes involved in epigenetic mechanisms in the published genome. BLAST searches suggest the presence of at least six HDACs, two HATs, five histone methyltransferases (HMTs), two histone demethylases (HDMs) and one MBD (supplementary material Table S2, Figs S1–S6).

A comparative methylome analysis of brain tissues between fourth instar solitary and gregarious *L. migratoria* (Wang et al., 2014) revealed lower and more fluctuating levels of CpG methylation in the coding regions of the genome compared with the genome as a whole. The ratios of observed/expected CpG levels show a bimodal distribution curve [as in *A. mellifera* and *S. gregaria* (Elango et al., 2009; Falckenhayn et al., 2013; Foret et al., 2009; Wang and Leung, 2009)] and suggest historical germline methylation, particularly in the coding regions of the genome. As in *S. gregaria* (Falckenhayn et al., 2013), repetitive elements are highly methylated; introns are more methylated than exons; and 90 genes are differentially methylated (at least four differentially methylated CpG sites) in gregarious versus solitary locusts, including genes involved in cytoskeleton formation. Wang and colleagues suggest that these genes might be involved in synaptic plasticity and, for the phase transition, point to a crucial role of microtubule dynamics control in locust brains.

### Histone modifications

The role of histone modifications has been less well studied in locusts. Using immunoassays for *S. gregaria*, we found that histone H3 contains phosphorylation (at serine 10 and 28, and threonine 3 and 11, respectively), tri-methylation and acetylation (both at lysine 9 and 27) (Boerjan et al., 2013). Preliminary data suggest that brains of gregarious *S. gregaria* contain more phosphorylated histone H3 than do those of solitary *S. gregaria*.

### ncRNA

In migratory locusts, burst expression of retro-elements has been observed in the egg stage, which is thought to be involved in locust development and has been proposed as a regulatory mechanism in phase transition (Guo et al., 2010). The involvement of small ncRNAs in *L. migratoria* phase transition was investigated by Wei et al., who compared small ncRNA abundance between the gregarious and solitary phase states (Wei et al., 2009). The two phases differed strongly in both length distribution and type of small RNAs. Gregarious animals had higher expression of small RNAs below 22 nucleotides, whereas the opposite was true for small RNAs above 22 nucleotides. Gregarious animals also had double the amount of miRNAs, whereas the solitary state expressed higher levels of endo-siRNAs and piRNA-like small RNAs. Moreover, miRNA-133 has been shown to inhibit behavioural aggregation by controlling dopamine synthesis in locusts (Yang et al., 2014). All

this is in strong support of an epigenetic basis for phase polymorphism.

In summary, increasing evidence points to a pivotal role of epigenetics in controlling phase transitions in locusts, yet definite evidence, i.e. more than correlational, is still lacking.

### Future research directions

Until recently, evidence for methylation in locusts relied on paper chromatography and photo-spectroscopy (Wyatt, 1951), mass spectrometry (Boerjan et al., 2011; Lechner et al., 2013) and methylation-specific restriction enzyme assays (Robinson et al., 2011). While these methods have their specific value, single base resolution methylome analyses allow detailed analysis of hypothetical changes in DNA methylation status between the solitary and gregarious phase (Robinson et al., 2011). Recently, these analyses have been performed in *L. migratoria* (Wang et al., 2014). Parts of a *Schistocerca* methylome have also been characterised (Falckenhayn et al., 2013) but, unfortunately, differences between phases were not investigated. Methylome studies to date have been based on whole organisms or brain tissue. However, it is likely that some epigenetic differences might be concealed by these approaches, as they might be specifically directed to particular brain regions or even cells (Bonasio, 2012).

### Pharmaceutical manipulation of global DNA methylation status

Strong evidence for a role of epigenetics in locust phase transition would be the experimental switch between phases in the treated animal and/or its offspring by altering parts of the epigenetic setting, be it DNA methylation, histone modification, ncRNAs, nucleosome positioning, or other. The epigenetic machinery can be experimentally manipulated, e.g. by blocking involved enzymes (such as Dnmts or MBDs), or by downregulating the synthesis of these enzymes. In recent years, it has even become possible to manipulate DNA methylation and histone modifications at specific sites (see below).

Gene regulation by DNA methylation is a complex matter (not to mention the intricate interactions with histones, their modifications and ncRNAs), and transitions from one state to another are usually characterised by a dynamic alteration in the methylation pattern. While some genes (or promoters) are being methylated, others are demethylated, and as such the total content of methylated cytosines could be more or less constant. For instance, most cancers are associated with a specific methylation pattern, where tumour suppressor genes are hypermethylated (i.e. inactivated), while oncogenes are hypomethylated (i.e. activated). It is evident that drugs or other experimental methods that aim at a general lower or higher level of DNA methylation are rather crude tools and will probably not be able to mimic such a delicate balance. Indeed, some cancer forms are associated with global hypomethylation, which also might promote metastasis (reviewed in Szyf, 2009). A complete demethylation is usually lethal, and Dnmts are often essential during development (e.g. Zwier et al., 2012).

Drugs like zebularine (Zhou et al., 2002), azacytidine, and decitabine (reviewed in Christman, 2002) are nucleoside analogues that prevent *de novo* and maintenance methylation by binding covalently to Dnmts when incorporated into DNA. As these drugs do not actively remove methyl groups, they are only effective where DNA replication takes place, limiting their usage to dividing cells.

Several other compounds that are not incorporated in DNA have been described as Dnmt inhibitors or DNA demethylating agents (reviewed in Szyf, 2009), but their mode of action is not yet understood. Mechanisms and enzymes that actively demethylate

DNA have been described in mammals and plants, the most prominent being TET (ten–eleven translocation) proteins (reviewed in Kohli and Zhang, 2013; Piccolo and Fisher, 2014; Wu and Zhang, 2014). Some of these demethylating pathways do not require DNA replication and would therefore be attractive to manipulate post-mitotic cells.

It should be noted that DNA methylation in insects is not necessarily associated with silencing of gene expression; in fact, DNA methylation in insects is highly correlated with steady gene expression and reduced variability in transcript levels (Foret et al., 2009; Lyko et al., 2010; Zemach et al., 2010).

#### Pharmaceutical manipulation of global histone modifications

Besides targeting DNA methylation, another promising avenue would be the manipulation of histone modifications. In humans, several classes of drugs have been developed for treating cancer and psychological disorders that interact with histone modifying enzymes such as HMT, HDM, HAT or HDAC (reviewed in Grayson et al., 2010; Szyf, 2009). Some of these enzymes are widely conserved, and could thus serve as targets for pharmaceuticals. Interestingly, HDAC inhibitors will not affect the whole genome, as both HDACs and HATs appear to be specific for certain sequences (reviewed in Szyf, 2009), which offers the opportunity to target a subset of genes. Some HDAC inhibitors also induce active DNA demethylation independent of DNA replication, which circumvents the disadvantages of drugs based on nucleoside analogues that are restricted to replicating DNA.

Similarly, HMT inhibitors will prevent the methylation of specific histones, which in turn prevent DNA methylation. In contrast, inhibition of histone demethylases would shift the balance between methylation and demethylation towards higher methylated histones. Histone (de)methylating enzymes are also specific, with several dozen specimens in several families described to date (reviewed in Højfeldt et al., 2013). So far, however, there are no reports of active demethylation processes in insects. While some drugs have been shown to work in humans and *Drosophila* alike (e.g. Greiner et al., 2005; but see Cherblanc et al., 2013), for some it remains to be evaluated whether they are also effective in other insects.

#### Analyses of DNA and histone modifications

Recent technologies allow mapping of different DNA modifications (methylation, hydroxymethylation, formylation, carboxylation) at single-base resolution (Booth et al., 2012; Raiber et al., 2012; Yu et al., 2012; Song et al., 2013), and we are only beginning to grasp the significance of these unusual nucleobases.

Various methods are available to map DNA methylation, each with their specific advantages and shortcomings (Bock et al., 2010; Harris et al., 2010; Laird, 2010; Mensaert et al., 2014; Nagarajan et al., 2013; Umer and Herceg, 2013). Wang and colleagues (Wang et al., 2014) used reduced representational bisulphite sequencing (RRBS) to map methylated CpG in the brains of solitary and gregarious locusts. This approach allows single-base resolution and absolute quantification of (hydroxy)methylation [albeit with imperfect quantification (Harris et al., 2010)], but is biased towards CpG-rich regions (Bock et al., 2010; Harris et al., 2010) and by the choice of the restriction enzyme used (Deng et al., 2009) and size selection of fragments (Bock et al., 2010). Both RRBS and whole-genome bisulphite sequencing (WGBS) may contain artifacts due to the bisulphite conversion process and subsequent PCR bias.

Studies of DNA modifications should be complemented by analyses of histone modifications where possible, e.g. by ChIP-Seq (chromatin immunoprecipitation followed by sequencing of the

immunoprecipitated DNA fragments) (Nagarajan et al., 2013; Rivera and Ren, 2013). A comparison of the epigenomes of the two phases between *L. migratoria* and *S. gregaria* will reveal the extent of common mechanisms as well as potentially specific modifications. Epigenetic changes that are shared between the two species should be prioritised in investigating their role in phase transition. It will be interesting to unravel how the different epigenetic pathways in these two species interact to bring about the phenomenon of phase transition.

#### Genome and epigenome editing

More specific tools to manipulate particular epigenetic signatures will have to be developed, e.g. ‘site-specific epimutagenesis’, as suggested previously (Bonasio, 2012). One strategy might be to target genes with a role in epigenetics using RNAi. Because of the robust systemic RNAi mechanism in locusts (Wynant et al., 2012), knockdown of target genes throughout the insect can easily be achieved by injecting dsRNA into the haemolymph. While RNAi has been successfully used for several years in locusts, it has one major drawback: it rarely induces a complete loss of function of the targeted gene. Innovative site-specific genomic engineering tools are currently being explored. For instance, zinc-finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs) or CRISPR-Cas (clustered regularly interspaced short palindromic repeat–CRISPR-associated proteins) could be used to specifically introduce mutations (indels, insertions and deletions) in target genes with roles in epigenetics, such as HATs and HDACs. CRISPR-Cas systems could even be used to edit several genes at once (Jao et al., 2013; Li et al., 2013; Wang et al., 2013). As HATs and HDACs are substrate specific, this would only affect a subset of histones. Alternatively, methylated DNA sequences of interest might be specifically excised and replaced by unmethylated DNA, or vice versa (Ramalingam et al., 2013).

Even more intriguing was the suggestion that methylating and demethylating enzymes as well as histone modifying enzymes (e.g. HMT, HAT) might be directed to specific DNA sequences of interest using transcription activator-like effector (TALE) or CRISPR-Cas (Bonasio, 2012; Gaj et al., 2013; Rivera and Ren, 2013). This would allow specific (de)methylation of DNA and histones, thereby enabling manipulation of epigenomes with high precision. This can be accomplished by using a catalytically inactive (or ‘dead’) version of Cas9, called dCas9, fused to an effector domain that will carry out its function (Sander and Joung, 2014). While the CRISPR-Cas system has already been shown to work in diverse organisms, such as bacteria, plants, vertebrates and insects, it has not yet been employed in locusts.

Recently, several groups succeeded in using TALE to selectively demethylate DNA (Maeder et al., 2013) and histones (Mendenhall et al., 2013). Konermann and colleagues (Konermann et al., 2013) were able to switch histone deacetylation and methylation on and off in a time- and space-specific manner using light-inducible transcriptional effectors (LITEs) in a combination of TALEs and optogenetic methods. This method is highly versatile, as it allows different DNA sequence recognition systems (e.g. zinc finger, TALE) to be combined with various ‘switches’ (e.g. light or chemical induced) and effectors (e.g. HDACs, HMTs). These manipulations would allow functional analyses of specific sites in the epigenome.

However, to date, locusts have not been genetically or epigenetically modified. One of the challenges will be to deliver the modified enzymes into cells of interest (Gaj et al., 2013). Interestingly, some ZFNs can be delivered directly as proteins across



the cell membrane (Gaj et al., 2012), but this is unlikely to work for the complex LITEs, which have been delivered by viral vectors (Konermann et al., 2013). In *D. melanogaster*, injection of cas9 – as DNA, RNA or protein – into the embryo allowed genetic manipulation that was transmitted to the offspring through the germline in 25–100% of cases, depending on the method and genes targeted (e.g. Bassett and Liu, 2014; Lee et al., 2014; Port et al., 2014). Given the ready use of RNAi in locusts (Wynant et al., 2012), it is possible that a similar approach will also work in locusts. If successful, this would open unprecedented opportunities for functional genomics and epigenomics, circumventing the drawbacks of RNAi and pharmaceutical approaches.

## Conclusions

Future research should aim to reveal the mechanisms of epigenetic control of locust phase transition. Additional insights are to be expected from comparisons with the Australian plague locust, *Chortoicetes terminifera*, where phase characteristics can change within 72 h in both directions and change abruptly between generations. Given the advent of new and exciting techniques that allow the specific manipulation of genes and proteins, and the analysis of single cells, these are excellent times to study the molecular mechanics of phase polyphenism in locusts.

## Acknowledgements

We apologise to those colleagues whose work we could not cite because of space limitations. We acknowledge the constructive suggestions of two anonymous reviewers that greatly helped to improve the article. We are grateful for the contributions of Liesbet Temmerman and Isabel Beets to the manuscript. Thanks are due to Tom Fayle for permission to use some of his images, to Roger Jonckers for taking care of our locusts and to Marijke Christiaens for assistance with the figures.

## Competing interests

The authors declare no competing financial interests.

## Author contributions

U.R.E., M.B.V.H., G.D., B.B., A.D.L. and L.S. drafted the manuscript. U.R.E. made the figures. M.B.V.H. made the table and supplementary data.

## Funding

This work was supported by the Research Foundation-Flanders (FWO) (Postdoctoral fellowship to B.B.), the Agency for Innovation by Science and Technology in Flanders (IWT) (U.R.E.) and a University of Leuven GOA grant (GOA/11/002).

## Supplementary material

Supplementary material available online at <http://jeb.biologists.org/lookup/suppl/doi:10.1242/jeb.107078/-DC1>

## References

- Achwal, C. W., Iyer, C. A. and Chandra, H. S. (1983). Immunochemical evidence for the presence of 5mC, 6mA and 7mG in human, *Drosophila* and mealworm DNA. *FEBS Lett.* **158**, 353–358.
- Amarasinghe, H. E., Clayton, C. I. and Mallon, E. B. (2014). Methylation and worker reproduction in the bumblebee (*Bombus terrestris*). *Proc. Biol. Sci.* **281**, 20132502.
- Anstey, M. L., Rogers, S. M., Ott, S. R., Burrows, M. and Simpson, S. J. (2009). Serotonin mediates behavioral gregarization underlying swarm formation in desert locusts. *Science* **323**, 627–630.
- Ayali, A., Perner, M. P. and Girardie, J. (1996). Comparative study of neuropeptides from the corpora cardiaca of solitary and gregarious *Locusta*. *Arch. Insect Biochem. Physiol.* **31**, 439–450.
- Ayali, A., Fuchs, E. and Kutsch, W. (2004). Neurophysiological studies of flight-related density-dependent phase characteristics in locusts. *Acta Biol. Hung.* **55**, 137–141.
- Badisco, L., Huybrechts, J., Simonet, G., Verlinden, H., Marchal, E., Huybrechts, R., Schoofs, L., De Loof, A. and Vanden Broeck, J. (2011a). Transcriptome analysis of the desert locust central nervous system: production and annotation of a *Schistocerca gregaria* EST database. *PLoS ONE* **6**, e17274.
- Badisco, L., Ott, S. R., Rogers, S. M., Matheson, T., Knapen, D., Vergauwen, L., Verlinden, H., Marchal, E., Sheehy, M. R., Burrows, M. et al. (2011b). Microarray-based transcriptomic analysis of differences between long-term gregarious and solitary desert locusts. *PLoS ONE* **6**, e28110.
- Baggerman, G., Huybrechts, J., Clynen, E., Hens, K., Harthoorn, L., Van der Horst, D., Poulos, C., De Loof, A. and Schoofs, L. (2002). New insights in Adipokinetic hormone (AKH) precursor processing in *Locusta migratoria* obtained by capillary liquid chromatography-tandem mass spectrometry. *Peptides* **23**, 635–644.
- Bannister, A. J. and Kouzarides, T. (2011). Regulation of chromatin by histone modifications. *Cell Res.* **21**, 381–395.
- Bassett, A. R. and Liu, J.-L. (2014). CRISPR/Cas9 and genome editing in *Drosophila*. *J. Genet. Genomics* **41**, 7–19.
- Beeler, S. M., Wong, G. T., Zheng, J. M., Bush, E. C., Remnant, E. J., Oldroyd, B. P. and Drewell, R. A. (2014). Whole-genome DNA methylation profile of the Jewel Wasp (*Nasonia vitripennis*). *G3 (Bethesda)* **4**, 383–388.
- Behura, S. K. and Whitfield, C. W. (2010). Correlated expression patterns of microRNA genes with age-dependent behavioural changes in honeybee. *Insect Mol. Biol.* **19**, 431–439.
- Bird, A. P. (1980). DNA methylation and the frequency of CpG in animal DNA. *Nucleic Acids Res.* **8**, 1499–1504.
- Bird, A. (2007). Perceptions of epigenetics. *Nature* **447**, 396–398.
- Blackburn, L. M., Ott, S. R., Matheson, T., Burrows, M. and Rogers, S. M. (2010). Motor neurone responses during a postural reflex in solitary and gregarious desert locusts. *J. Insect Physiol.* **56**, 902–910.
- Bock, C., Tomazou, E. M., Brinkman, A. B., Müller, F., Simmer, F., Gu, H., Jäger, N., Gnirke, A., Stunnenberg, H. G. and Meissner, A. (2010). Quantitative comparison of genome-wide DNA methylation mapping technologies. *Nat. Biotechnol.* **28**, 1106–1114.
- Boerjan, B., Verleyen, P., Huybrechts, J., Schoofs, L. and De Loof, A. (2010). In search for a common denominator for the diverse functions of arthropod corazonin: a role in the physiology of stress? *Gen. Comp. Endocrinol.* **166**, 222–233.
- Boerjan, B., Sas, F., Ernst, U. R., Tobback, J., Lemièr, F., Vandegheuchte, M. B., Janssen, C. R., Badisco, L., Marchal, E., Verlinden, H. et al. (2011). Locust phase polyphenism: Does epigenetic precede endocrine regulation? *Gen. Comp. Endocrinol.* **173**, 120–128.
- Boerjan, B., Van Eynde, A., Bollen, M., De Loof, A. and Schoofs, L. (2013). Can epigenetics explain the swarm? Metaleptea – Special conference issue (11th International Congress of Orthopterology, Kummig, 11–15th August 2013). p. 173–175.
- Bonasio, R. (2012). Emerging topics in epigenetics: ants, brains, and noncoding RNAs. *Ann. N. Y. Acad. Sci.* **1260**, 14–23.
- Bonasio, R. (2014). The role of chromatin and epigenetics in the polyphenisms of ant castes. *Brief. Funct. Genomics* **13**, 235–245.
- Bonasio, R. (2015). The expanding epigenetic landscape of non-model organisms. *J. Exp. Biol.* **218**, 114–122.
- Bonasio, R., Zhang, G., Ye, C., Mutti, N. S., Fang, X., Qin, N., Donahue, G., Yang, P., Li, Q., Li, C. et al. (2010). Genomic comparison of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Science* **329**, 1068–1071.
- Bonasio, R., Li, Q., Lian, J., Mutti, N. S., Jin, L., Zhao, H., Zhang, P., Wen, P., Xiang, H., Ding, Y. et al. (2012). Genome-wide and caste-specific DNA methylomes of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Curr. Biol.* **22**, 1755–1764.
- Booth, M. J., Branco, M. R., Ficiz, G., Oxley, D., Krueger, F., Reik, W. and Balasubramanian, S. (2012). Quantitative sequencing of 5-methylcytosine and 5-hydroxymethylcytosine at single-base resolution. *Science* **336**, 934–937.
- Brakefield, P. M. and Frankino, W. A. (2009). Polyphenisms in Lepidoptera: multidisciplinary approaches to studies of evolution. In *Phenotypic Plasticity of Insects: Mechanisms and Consequences* (ed. D. Whitman and T. N. Ananthakrishnan), pp. 337–368. Enfield, NH: Science Publishers.
- Breuer, M., Hoste, B. and De Loof, A. (2003). The endocrine control of phase transition: some new aspects. *Physiol. Entomol.* **28**, 3–10.
- Burggren, W. W. (2015). Dynamics of epigenetic phenomena: intergenerational and intragenerational phenotype ‘washout’. *J. Exp. Biol.* **218**, 80–87.
- Burrows, M., Rogers, S. M. and Ott, S. R. (2011). Epigenetic remodelling of brain, body and behaviour during phase change in locusts. *Neural Syst. Circuits* **1**, 11.
- Chen, S., Yang, P., Jiang, F., Wei, Y., Ma, Z. and Kang, L. (2010). De novo analysis of transcriptome dynamics in the migratory locust during the development of phase traits. *PLoS ONE* **5**, e15633.
- Cherblanc, F. L., Chapman, K. L., Brown, R. and Fuchter, M. J. (2013). Chaetocin is a nonspecific inhibitor of histone lysine methyltransferases. *Nat. Chem. Biol.* **9**, 136–137.
- Christman, J. K. (2002). 5-Azacytidine and 5-aza-2'-deoxycytidine as inhibitors of DNA methylation: mechanistic studies and their implications for cancer therapy. *Oncogene* **21**, 5483–5495.
- Claeys, I., Simonet, G., Breugelmans, B., Van Soest, S., Franssens, V., Sas, F., De Loof, A. and Vanden Broeck, J. (2005). Quantitative real-time RT-PCR analysis in desert locusts reveals phase dependent differences in neuropeptide transcript levels. *Insect Mol. Biol.* **14**, 415–422.
- Claeys, I., Breugelmans, B., Simonet, G., Franssens, V., Van Soest, S. and Broeck, J. V. (2006a). Regulation of *Schistocerca gregaria* neuropeptide transcript levels by juvenile hormone and 20-hydroxyecdysone. *Arch. Insect Biochem. Physiol.* **62**, 107–115.
- Claeys, I., Breugelmans, B., Simonet, G., Van Soest, S., Sas, F., De Loof, A. and Vanden Broeck, J. (2006b). Neuropeptide transcripts as molecular markers in the process of desert locust (*Schistocerca gregaria*) phase transition. *Biochem. Biophys. Res. Commun.* **341**, 599–606.
- Clynen, E., Stubbe, D., De Loof, A. and Schoofs, L. (2002). Peptide differential display: a novel approach for phase transition in locusts. *Comp. Biochem. Physiol.* **132B**, 107–115.

- Cullen, D. A., Sword, G. A., Dodgson, T. and Simpson, S. J. (2010). Behavioural phase change in the Australian plague locust, *Chortoicetes terminifera*, is triggered by tactile stimulation of the antennae. *J. Insect Physiol.* **56**, 937-942.
- Deng, J., Shoemaker, R., Xie, B., Gore, A., LeProust, E. M., Antosiewicz-Bourget, J., Egli, D., Maherali, N., Park, I. H., Yu, J. et al. (2009). Targeted bisulfite sequencing reveals changes in DNA methylation associated with nuclear reprogramming. *Nat. Biotechnol.* **27**, 353-360.
- Dickman, M. J., Kucharski, R., Maleszka, R. and Hurd, P. J. (2013). Extensive histone post-translational modification in honey bees. *Insect Biochem. Mol. Biol.* **43**, 125-137.
- Duncan, B. K. and Miller, J. H. (1980). Mutagenic deamination of cytosine residues in DNA. *Nature* **287**, 560-561.
- Elango, N., Hunt, B. G., Goodisman, M. A. D. and Yi, S. V. (2009). DNA methylation is widespread and associated with differential gene expression in castes of the honeybee, *Apis mellifera*. *Proc. Natl. Acad. Sci. USA* **106**, 11206-11211.
- Ellis, P. E. (1953). Social aggregation and gregarious behaviour in hoppers of *Locusta migratoria migratorioides* (R. and F.). *Behaviour* **5**, 225-259.
- Ellis, P. E. (1959). Some factors influencing phase characters in the nymphs of the locust, *Locusta migratoria migratorioides* (R. and F.). *Insectes Soc.* **6**, 21-39.
- Ellis, P. E. (1962). The behaviour of locusts in relation to phases and species. *Colloq. Int. C.N.R.S.* **114**, 123-143.
- Falckenhayn, C., Boerjan, B., Raddatz, G., Frohme, M., Schoofs, L. and Lyko, F. (2013). Characterization of genome methylation patterns in the desert locust *Schistocerca gregaria*. *J. Exp. Biol.* **216**, 1423-1429.
- Ferenz, H. J. (1990). Locust pheromones- basic and applied aspects. *Boletín de Sanidad Vegetal, Fuere de Serie* **20**, 29-37. (Also as Proceedings of the 5th International Meeting of the Orthopterists' Society 1989).
- Filion, G. J., van Bommel, J. G., Braunschweig, U., Talhout, W., Kind, J., Ward, L. D., Brugman, W., de Castro, I. J., Kerkhoven, R. M., Bussemaker, H. J. et al. (2010). Systematic protein location mapping reveals five principal chromatin types in *Drosophila* cells. *Cell* **143**, 212-224.
- Foret, S., Kucharski, R., Pittelkow, Y., Lockett, G. A. and Maleszka, R. (2009). Epigenetic regulation of the honey bee transcriptome: unravelling the nature of methylated genes. *BMC Genomics* **10**, 472.
- Fuchs, E., Kutsch, W. and Ayali, A. (2003). Neural correlates to flight-related density-dependent phase characteristics in locusts. *J. Neurobiol.* **57**, 152-162.
- Gaj, T., Guo, J., Kato, Y., Sirk, S. J. and Barbas, C. F., III (2012). Targeted gene knockout by direct delivery of zinc-finger nuclease proteins. *Nat. Methods* **9**, 805-807.
- Gaj, T., Gersbach, C. A. and Barbas, C. F., 3rd (2013). ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. *Trends Biotechnol.* **31**, 397-405.
- Glastad, K. M., Hunt, B. G., Yi, S. V. and Goodisman, M. A. (2011). DNA methylation in insects: on the brink of the epigenomic era. *Insect Mol. Biol.* **20**, 553-565.
- Gnyoszka, A., Jastrzebski, Z. and Flis, S. (2013). DNA methyltransferase inhibitors and their emerging role in epigenetic therapy of cancer. *Anticancer Res.* **33**, 2989-2996.
- Grayson, D. R., Kundakovic, M. and Sharma, R. P. (2010). Is there a future for histone deacetylase inhibitors in the pharmacotherapy of psychiatric disorders? *Mol. Pharmacol.* **77**, 126-135.
- Greenberg, J. K., Xia, J., Zhou, X., Thatcher, S. R., Gu, X., Ament, S. A., Newman, T. C., Green, P. J., Zhang, W., Robinson, G. E. et al. (2012). Behavioral plasticity in honey bees is associated with differences in brain microRNA transcriptome. *Genes Brain Behav.* **11**, 660-670.
- Greiner, D., Bonaldi, T., Eskeland, R., Roemer, E. and Imhof, A. (2005). Identification of a specific inhibitor of the histone methyltransferase SU(VAR)3-9. *Nat. Chem. Biol.* **1**, 143-145.
- Guo, W., Wang, X. H., Zhao, D. J., Yang, P. C. and Kang, L. (2010). Molecular cloning and temporal-spatial expression of I element in gregarious and solitary locusts. *J. Insect Physiol.* **56**, 943-948.
- Guo, W., Wang, X., Ma, Z., Xue, L., Han, J., Yu, D. and Kang, L. (2011). CSP and takeout genes modulate the switch between attraction and repulsion during behavioral phase change in the migratory locust. *PLoS Genet.* **7**, e1001291.
- Guo, X., Ma, Z. and Kang, L. (2013a). Serotonin enhances solitariness in phase transition of the migratory locust. *Front. Behav. Neurosci.* **7**, 129.
- Guo, X., Su, S., Skogerboe, G., Dai, S., Li, W., Li, Z., Liu, F., Ni, R., Guo, Y., Chen, S. et al. (2013b). Recipe for a busy bee: microRNAs in Honey Bee caste determination. *PLoS ONE* **8**, e81661.
- Haig, D. (2004). The (dual) origin of epigenetics. *Cold Spring Harb. Symp. Quant. Biol.* **69**, 67-70.
- Halfmann, R. and Lindquist, S. (2010). Epigenetics in the extreme: prions and the inheritance of environmentally acquired traits. *Science* **330**, 629-632.
- Halfmann, R., Alberti, S. and Lindquist, S. (2010). Prions, protein homeostasis, and phenotypic diversity. *Trends Cell Biol.* **20**, 125-133.
- Halfmann, R., Jarosz, D. F., Jones, S. K., Chang, A., Lancaster, A. K. and Lindquist, S. (2012). Prions are a common mechanism for phenotypic inheritance in wild yeasts. *Nature* **482**, 363-368.
- Harris, R. A., Wang, T., Coarfa, C., Nagarajan, R. P., Hong, C., Downey, S. L., Johnson, B. E., Fouse, S. D., Delaney, A., Zhao, Y. et al. (2010). Comparison of sequencing-based methods to profile DNA methylation and identification of monoallelic epigenetic modifications. *Nat. Biotechnol.* **28**, 1097-1105.
- Hartfelder, K. and Emlen, D. J. (2012). Endocrine control of insect polyphenism. In *Insect Endocrinology* (ed. L. I. Gilbert), pp. 464-522. London; Waltham, MA: Elsevier/Academic Press.
- Hasegawa, E. and Tanaka, S. (1994). Genetic control of albinism and the role of juvenile hormone in pigmentation in *Locusta migratoria* (Orthoptera, Acrididae). *Jap. J. Entomol.* **62**, 315-324.
- Hassanali, A., Njagi, P. G. and Bashir, M. O. (2005). Chemical ecology of locusts and related acridids. *Annu. Rev. Entomol.* **50**, 223-245.
- Hatle, J. D. and Spring, J. H. (1999). Tests of potential adipokinetic hormone precursor related peptide (APRP) functions: lack of responses. *Arch. Insect Biochem. Physiol.* **42**, 163-166.
- Heifetz, Y., Voet, H. and Applebaum, S. W. (1996). Factors affecting behavioral phase transition in the desert locust, *Schistocerca gregaria* (Forskål) (Orthoptera: Acrididae). *J. Chem. Ecol.* **22**, 1717-1734.
- Heifetz, Y., Boekhoff, I., Breer, H. and Applebaum, S. W. (1997). Cuticular hydrocarbons control behavioural phase transition in *Schistocerca gregaria* nymphs and elicit biochemical responses in antennae. *Insect Biochem. Mol. Biol.* **27**, 563-568.
- Heifetz, Y., Miloslavski, I., Aizenshtat, Z. and Applebaum, S. W. (1998). Cuticular surface hydrocarbons of Desert Locust nymphs, *Schistocerca gregaria*, and their effect on phase behavior. *J. Chem. Ecol.* **24**, 1033-1047.
- Herb, B. R., Wolschin, F., Hansen, K. D., Aryee, M. J., Langmead, B., Irizarry, R., Amdam, G. V. and Feinberg, A. P. (2012). Reversible switching between epigenetic states in honeybee behavioral subcastes. *Nat. Neurosci.* **15**, 1371-1373.
- Højfeldt, J. W., Agger, K. and Helin, K. (2013). Histone lysine demethylases as targets for anticancer therapy. *Nat. Rev. Drug Discov.* **12**, 917-930.
- Hoste, B., Simpson, S. J., Tanaka, S., Zhu, D., De Loof, A. and Breuer, M. (2002). Effects of [His(7)]-corazonin on the phase state of isolated-reared (solitary) desert locusts, *Schistocerca gregaria*. *J. Insect Physiol.* **48**, 981-990.
- Hoste, B., Simpson, S. J., De Loof, A. and Breuer, M. (2003). Behavioural differences in *Locusta migratoria* associated with albinism and their relation to [His(7)]-corazonin. *Physiol. Entomol.* **28**, 32-38.
- Hunt, B. G., Glastad, K. M., Yi, S. V. and Goodisman, M. A. (2013a). The function of intragenic DNA methylation: insights from insect epigenomes. *Integr. Comp. Biol.* **53**, 319-328.
- Hunt, B. G., Glastad, K. M., Yi, S. V. and Goodisman, M. A. (2013b). Patterning and regulatory associations of DNA methylation are mirrored by histone modifications in insects. *Genome Biol. Evol.* **5**, 591-598.
- Islam, M. S. (2013). Behavioural interpretations of the HPLC peaks derived from egg-pod foam extracts of the Desert Locust *Schistocerca gregaria* (Forskål). *Annual Review & Research in Biology* **3**, 475-491.
- Jablonka, E. and Lamb, M. J. (2002). The changing concept of epigenetics. *Ann. N. Y. Acad. Sci.* **981**, 82-96.
- Jablonka, E. and Raz, G. (2009). Transgenerational epigenetic inheritance: prevalence, mechanisms, and implications for the study of heredity and evolution. *Q. Rev. Biol.* **84**, 131-176.
- Jacquier, A. (2009). The complex eukaryotic transcriptome: unexpected pervasive transcription and novel small RNAs. *Nat. Rev. Genet.* **10**, 833-844.
- Jao, L. E., Wente, S. R. and Chen, W. (2013). Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system. *Proc. Natl. Acad. Sci. USA* **110**, 13904-13909.
- Jenuwein, T. and Allis, C. D. (2001). Translating the histone code. *Science* **293**, 1074-1080.
- Jiang, F., Yang, M., Guo, W., Wang, X. and Kang, L. (2012). Large-scale transcriptome analysis of retroelements in the migratory locust, *Locusta migratoria*. *PLoS ONE* **7**, e40532.
- Kang, L., Chen, X., Zhou, Y., Liu, B., Zheng, W., Li, R., Wang, J. and Yu, J. (2004). The analysis of large-scale gene expression correlated to the phase changes of the migratory locust. *Proc. Natl. Acad. Sci. USA* **101**, 17611-17615.
- Kharchenko, P. V., Alekseyenko, A. A., Schwartz, Y. B., Minoda, A., Riddle, N. C., Ernst, J., Sabo, P. J., Larschan, E., Gorchakov, A. A., Gu, T. et al. (2011). Comprehensive analysis of the chromatin landscape in *Drosophila melanogaster*. *Nature* **471**, 480-485.
- Kijimoto, T., Pespini, M., Beckers, O. and Moczek, A. P. (2013). Beetle horns and horned beetles: emerging models in developmental evolution and ecology. *Wiley Interdiscip. Rev. Dev. Biol.* **2**, 405-418.
- Klass, M., Nguyen, P. N. and Dechavigny, A. (1983). Age-correlated changes in the DNA template in the nematode *Caenorhabditis elegans*. *Mech. Ageing Dev.* **22**, 253-263.
- Kohli, R. M. and Zhang, Y. (2013). TET enzymes, TDG and the dynamics of DNA demethylation. *Nature* **502**, 472-479.
- Konermann, S., Brigham, M. D., Trevino, A. E., Hsu, P. D., Heidenreich, M., Cong, L., Platt, R. J., Scott, D. A., Church, G. M. and Zhang, F. (2013). Optical control of mammalian endogenous transcription and epigenetic states. *Nature* **500**, 472-476.
- Krauss, V., Eisenhardt, C. and Unger, T. (2009). The genome of the stick insect *Medauroidea extradentata* is strongly methylated within genes and repetitive DNA. *PLoS ONE* **4**, e7223.
- Kucharski, R., Maleszka, J., Foret, S. and Maleszka, R. (2008). Nutritional control of reproductive status in honeybees via DNA methylation. *Science* **319**, 1827-1830.
- Laird, P. W. (2010). Principles and challenges of genomewide DNA methylation analysis. *Nat. Rev. Genet.* **11**, 191-203.
- Lechner, M., Marz, M., Ihling, C., Sinz, A., Stadler, P. F. and Krauss, V. (2013). The correlation of genome size and DNA methylation rate in metazoans. *Theory Biosci.* **132**, 47-60.
- Lee, J. S., Kwak, S. J., Kim, J., Kim, A. K., Noh, H. M., Kim, J. S. and Yu, K. (2014). RNA-guided genome editing in *drosophila* with the purified Cas9 protein. *G3 (Bethesda)* **4**, 1291-1295.



- Lenz, E. M., Hägele, B. F., Wilson, I. D. and Simpson, S. J. (2001). High resolution <sup>1</sup>H NMR spectroscopic studies of the composition of the haemolymph of crowd- and solitary-reared nymphs of the desert locust, *Schistocerca gregaria*. *Insect Biochem. Mol. Biol.* **32**, 51-56.
- Leo Lester, R., Grach, C., Paul Pener, M. and Simpson, S. J. (2005). Stimuli inducing gregarious colouration and behaviour in nymphs of *Schistocerca gregaria*. *J. Insect Physiol.* **51**, 737-747.
- Li, W., Teng, F., Li, T. and Zhou, Q. (2013). Simultaneous generation and germline transmission of multiple gene mutations in rat using CRISPR-Cas systems. *Nat. Biotechnol.* **31**, 684-686.
- Li-Byarlay, H., Li, Y., Stroud, H., Feng, S., Newman, T. C., Kaneda, M., Hou, K. K., Worley, K. C., Elsik, C. G., Wickline, S. A. et al. (2013). RNA interference knockdown of DNA methyl-transferase 3 affects gene alternative splicing in the honey bee. *Proc. Natl. Acad. Sci. USA* **110**, 12750-12755.
- Liu, F., Peng, W., Li, Z., Li, W., Li, L., Pan, J., Zhang, S., Miao, Y., Chen, S. and Su, S. (2012). Next-generation small RNA sequencing for microRNAs profiling in *Apis mellifera*: comparison between nurses and foragers. *Insect Mol. Biol.* **21**, 297-303.
- Lo, N., Li, B. and Ujvari, B. (2012). DNA methylation in the termite *Coptotermes lacteus*. *Insectes Soc.* **59**, 257-261.
- Lockett, G. A., Kucharski, R. and Maleszka, R. (2012). DNA methylation changes elicited by social stimuli in the brains of worker honey bees. *Genes Brain Behav.* **11**, 235-242.
- Lyko, F. and Maleszka, R. (2011). Insects as innovative models for functional studies of DNA methylation. *Trends Genet.* **27**, 127-131.
- Lyko, F., Ramsahoye, B. H. and Jaenisch, R. (2000). DNA methylation in *Drosophila melanogaster*. *Nature* **408**, 538-540.
- Lyko, F., Foret, S., Kucharski, R., Wolf, S., Falckenhayn, C. and Maleszka, R. (2010). The honey bee epigenomes: differential methylation of brain DNA in queens and workers. *PLoS Biol.* **8**, e1000506.
- Ma, Z., Guo, W., Guo, X., Wang, X. and Kang, L. (2011). Modulation of behavioral phase changes of the migratory locust by the catecholamine metabolic pathway. *Proc. Natl. Acad. Sci. USA* **108**, 3882-3887.
- Maeder, M. L., Angstman, J. F., Richardson, M. E., Linder, S. J., Cascio, V. M., Tsai, S. Q., Ho, Q. H., Sander, J. D., Reyon, D., Bernstein, B. E. et al. (2013). Targeted DNA demethylation and activation of endogenous genes using programmable TALE-TET1 fusion proteins. *Nat. Biotechnol.* **31**, 1137-1142.
- Maeno, K. and Tanaka, S. (2004). Hormonal control of phase-related changes in the number of antennal sensilla in the desert locust, *Schistocerca gregaria*: possible involvement of [His7]-corazonin. *J. Insect Physiol.* **50**, 855-865.
- Maeno, K. and Tanaka, S. (2008). Maternal effects on progeny size, number and body color in the desert locust, *Schistocerca gregaria*: density- and reproductive cycle-dependent variation. *J. Insect Physiol.* **54**, 1072-1080.
- Maeno, K. and Tanaka, S. (2009). Is juvenile hormone involved in the maternal regulation of egg size and progeny characteristics in the desert locust? *J. Insect Physiol.* **55**, 1021-1028.
- Maeno, K., Gotoh, T. and Tanaka, S. (2004). Phase-related morphological changes induced by [His7]-corazonin in two species of locusts, *Schistocerca gregaria* and *Locusta migratoria* (Orthoptera: Acrididae). *Bull. Entomol. Res.* **94**, 349-357.
- Maeno, K., Tanaka, S. and Harano, K. (2011). Tactile stimuli perceived by the antennae cause the isolated females to produce gregarious offspring in the desert locust, *Schistocerca gregaria*. *J. Insect Physiol.* **57**, 74-82.
- Mendenhall, E. M., Williamson, K. E., Reyon, D., Zou, J. Y., Ram, O., Joung, J. K. and Bernstein, B. E. (2013). Locus-specific editing of histone modifications at endogenous enhancers. *Nat. Biotechnol.* **31**, 1133-1136.
- Mensaert, K., Denil, S., Trooskens, G., Van Crielinge, W., Thas, O. and De Meyer, T. (2014). Next-generation technologies and data analytical approaches for epigenomics. *Environ. Mol. Mutagen.* **55**, 155-170.
- Miller, G. A., Islam, M. S., Claridge, T. D., Dodgson, T. and Simpson, S. J. (2008). Swarm formation in the desert locust *Schistocerca gregaria*: isolation and NMR analysis of the primary maternal gregarizing agent. *J. Exp. Biol.* **211**, 370-376.
- Moazed, D. (2009). Small RNAs in transcriptional gene silencing and genome defence. *Nature* **457**, 413-420.
- Moczek, A. P. (2010). Phenotypic plasticity and diversity in insects. *Philos. Trans. R. Soc. B* **365**, 593-603.
- Nagarajan, R. P., Fouse, S. D., Bell, R. J. and Costello, J. F. (2013). Methods for cancer epigenome analysis. *Adv. Exp. Med. Biol.* **754**, 313-338.
- Nanty, L., Carbajosa, G., Heap, G. A., Ratnieks, F., van Heel, D. A., Down, T. A. and Rakyan, V. K. (2011). Comparative methylomics reveals gene-body H3K36me3 in *Drosophila* predicts DNA methylation and CpG landscapes in other invertebrates. *Genome Res.* **21**, 1841-1850.
- Navarro-Martin, L., Viñas, J., Ribas, L., Díaz, N., Gutiérrez, A., Di Croce, L. and Piferrer, F. (2011). DNA methylation of the gonadal aromatase (*cyp19a*) promoter is involved in temperature-dependent sex ratio shifts in the European sea bass. *PLoS Genet.* **7**, e1002447.
- Nègre, N., Brown, C. D., Ma, L., Bristow, C. A., Miller, S. W., Wagner, U., Kheradpour, P., Eaton, M. L., Loriaux, P., Sealfon, R. et al. (2011). A cis-regulatory map of the *Drosophila* genome. *Nature* **471**, 527-531.
- Norris, M. J. (1950). Reproduction in the African Migratory locust (*Locusta migratoria migratorioides* R. & F.) in relation to density and phase. *Anti-Locust Bulletin* **6**, 1-48.
- Norris, M. J. (1952). Reproduction in the desert locust (*Schistocerca gregaria* Forsk.) in relation to density and phase. *Anti-Locust Bulletin* **13**, 1-51.
- Ogawa, K. and Miura, T. (2014). Aphid polyphenisms: trans-generational developmental regulation through viviparity. *Front. Physiol.* **5**, 1.
- Ott, S. R. and Rogers, S. M. (2010). Gregarious desert locusts have substantially larger brains with altered proportions compared with the solitary phase. *Proc. Biol. Sci.* **277**, 3087-3096.
- Ott, S. R., Verlinden, H., Rogers, S. M., Brighton, C. H., Quah, P. S., Vleugels, R. K., Verdonck, R. and Vanden Broeck, J. (2012). Critical role for protein kinase A in the acquisition of gregarious behavior in the desert locust. *Proc. Natl. Acad. Sci. USA* **109**, E381-E387.
- Pauli, A., Rinn, J. L. and Schier, A. F. (2011). Non-coding RNAs as regulators of embryogenesis. *Nat. Rev. Genet.* **12**, 136-149.
- Pener, M. (1991). Locust phase polymorphism and its endocrine relations. *Adv. In Insect Phys.* **23**, 1-79.
- Pener, M. and Simpson, S. (2009). Locust phase polyphenism: an update. *Adv. In Insect Phys.* **36**, 1-272.
- Pener, M. P. and Yerushalmi, Y. (1998). The physiology of locust phase polymorphism: an update. *J. Insect Physiol.* **44**, 365-377.
- Piccolo, F. M. and Fisher, A. G. (2014). Getting rid of DNA methylation. *Trends Cell Biol.* **24**, 136-143.
- Poirier, S., Samami, S., Mamarbachi, M., Demers, A., Chang, T. Y., Vance, D. E., Hatch, G. M. and Mayer, G. (2014). The epigenetic drug 5-azacytidine interferes with cholesterol and lipid metabolism. *J. Biol. Chem.* **289**, 18736-18751.
- Port, F., Chen, H. M., Lee, T. and Bullock, S. L. (2014). Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **111**, E2967-E2976.
- Raddatz, G., Guzzardo, P. M., Olova, N., Fantappiè, M. R., Rampp, M., Schaefer, M., Reik, W., Hannon, G. J. and Lyko, F. (2013). Dnmt2-dependent methylomes lack defined DNA methylation patterns. *Proc. Natl. Acad. Sci. USA* **110**, 8627-8631.
- Rahman, M. M., Bosch, L. V., Baggerman, G., Clynen, E., Hens, K., Hoste, B., Meylaers, K., Vercammen, T., Schoofs, L., De Loof, A. et al. (2002). Search for peptidic molecular markers in hemolymph of crowd-(gregarious) and isolated-reared (solitary) desert locusts, *Schistocerca gregaria*. *Peptides* **23**, 1907-1914.
- Rahman, M., Baggerman, G., Begum, M., De Loof, A. and Breuer, M. (2003a). Purification, isolation and search for possible functions of a phase-related 6080-Da peptide from the haemolymph of the desert locust, *Schistocerca gregaria*. *Physiol. Entomol.* **28**, 39-45.
- Rahman, M. M., Vandingenen, A., Begum, M., Breuer, M., De Loof, A. and Huybrechts, R. (2003b). Search for phase specific genes in the brain of desert locust, *Schistocerca gregaria* (Orthoptera: Acrididae) by differential display polymerase chain reaction. *Comp. Biochem. Physiol.* **135A**, 221-228.
- Raiber, E. A., Beraldi, D., Ficiz, G., Burgess, H. E., Branco, M. R., Murat, P., Oxley, D., Booth, M. J., Reik, W. and Balasubramanian, S. (2012). Genome-wide distribution of 5-formylcytosine in embryonic stem cells is associated with transcription and depends on thymine DNA glycosylase. *Genome Biol.* **13**, R69.
- Ramalingam, S., Annaluru, N. and Chandrasegaran, S. (2013). A CRISPR way to engineer the human genome. *Genome Biol.* **14**, 107.
- Riggs, A. D., Martienssen, R. A. and Russo, V. E. A. 1996. Introduction. In *Epigenetic Mechanisms of Gene Regulation* (ed. V. E. A. Russo, R. A. Martienssen and A. D. Riggs), pp. 1. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Rivera, C. M. and Ren, B. (2013). Mapping human epigenomes. *Cell* **155**, 39-55.
- Robinson, K. L., Tohidi-Esfahani, D., Lo, N., Simpson, S. J. and Sword, G. A. (2011). Evidence for widespread genomic methylation in the migratory locust, *Locusta migratoria* (Orthoptera: Acrididae). *PLoS ONE* **6**, e28167.
- Roessingh, P., Simpson, S. and James, S. (1993). Analysis of phase-related changes in behaviour of desert locust nymphs. *Proc. Biol. Sci.* **252**, 43-49.
- Roessingh, P., Bouaichi, A. and Simpson, S. J. (1998). Effects of sensory stimuli on the behavioural phase state of the desert locust, *Schistocerca gregaria*. *J. Insect Physiol.* **44**, 883-893.
- Rogers, S. M., Matheson, T., Despland, E., Dodgson, T., Burrows, M. and Simpson, S. J. (2003). Mechanosensory-induced behavioural gregarization in the desert locust *Schistocerca gregaria*. *J. Exp. Biol.* **206**, 3991-4002.
- Rogers, S. M., Matheson, T., Sasaki, K., Kendrick, K., Simpson, S. J. and Burrows, M. (2004). Substantial changes in central nervous system neurotransmitters and neuromodulators accompany phase change in the locust. *J. Exp. Biol.* **207**, 3603-3617.
- Sander, J. D. and Joung, J. K. (2014). CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat. Biotechnol.* **32**, 347-355.
- Sarkar, S., Rao, S. R. V., Gupta, V. S. and Hendre, R. R. (1992). 5-Methylcytosine content in *Gryllotalpa fossor* (Orthoptera). *Genome* **35**, 163-166.
- Sas, F., Begum, M., Vandersmissen, T., Geens, M., Claeys, I., Van Soest, S., Huybrechts, J., Huybrechts, R. and De Loof, A. (2007). Development of a real-time PCR assay for measurement of yellow protein mRNA transcription in the desert locust *Schistocerca gregaria*: a basis for isolation of a peptidergic regulatory factor. *Peptides* **28**, 38-43.
- Simões, P. M., Niven, J. E. and Ott, S. R. (2013). Phenotypic transformation affects associative learning in the desert locust. *Curr. Biol.* **23**, 2407-2412.
- Simola, D. F., Ye, C., Mutti, N. S., Dolezal, K., Bonasio, R., Liebig, J., Reinberg, D. and Berger, S. L. (2013). A chromatin link to caste identity in the carpenter ant *Camponotus floridanus*. *Genome Res.* **23**, 486-496.
- Simonet, G., Claeys, I., Breugelmans, B., Van Soest, S., De Loof, A. and Vanden Broeck, J. (2004). Transcript profiling of pacifastin-like peptide precursors in crowd- and isolated-reared desert locusts. *Biochem. Biophys. Res. Commun.* **317**, 565-569.
- Simonet, G., Breugelmans, B., Proost, P., Claeys, I., Van Damme, J., De Loof, A. and Vanden Broeck, J. (2005). Characterization of two novel pacifastin-like peptide precursor isoforms in the desert locust (*Schistocerca gregaria*): cDNA cloning, functional analysis and real-time RT-PCR gene expression studies. *Biochem. J.* **388**, 281-289.



- Simpson, S. J. and Miller, G. A. (2007). Maternal effects on phase characteristics in the desert locust, *Schistocerca gregaria*: a review of current understanding. *J. Insect Physiol.* **53**, 869–876.
- Simpson, V. J., Johnson, T. E. and Hammen, R. F. (1986). *Caenorhabditis elegans* DNA does not contain 5-methylcytosine at any time during development or aging. *Nucleic Acids Res.* **14**, 6711–6719.
- Simpson, S. J., McCaffery, A. R. and Hägele, B. F. (1999). A behavioural analysis of phase change in the desert locust. *Biol. Rev. Camb. Philos. Soc.* **74**, 461–480.
- Simpson, S. J., Despland, E., Hägele, B. F. and Doddson, T. (2001). Gregarious behavior in desert locusts is evoked by touching their back legs. *Proc. Natl. Acad. Sci. USA* **98**, 3895–3897.
- Simpson, S. J., Sword, G. A. and Lo, N. (2011). Polyphenism in insects. *Curr. Biol.* **21**, R738–R749.
- Slotkin, R. K. and Martienssen, R. (2007). Transposable elements and the epigenetic regulation of the genome. *Nat. Rev. Genet.* **8**, 272–285.
- Smith, C. R., Mutti, N. S., Jasper, W. C., Naidu, A., Smith, C. D. and Gadau, J. (2012). Patterns of DNA methylation in development, division of labor and hybridization in an ant with genetic caste determination. *PLoS ONE* **7**, e42433.
- Song, C. X., Szulwach, K. E., Dai, Q., Fu, Y., Mao, S. Q., Lin, L., Street, C., Li, Y., Poidevin, M., Wu, H. et al. (2013). Genome-wide profiling of 5-formylcytosine reveals its roles in epigenetic priming. *Cell* **153**, 678–691.
- Strahl, B. D. and Allis, C. D. (2000). The language of covalent histone modifications. *Nature* **403**, 41–45.
- Suganuma, T. and Workman, J. L. (2011). Signals and combinatorial functions of histone modifications. *Annu. Rev. Biochem.* **80**, 473–499.
- Szyf, M. (2009). Epigenetics, DNA methylation, and chromatin modifying drugs. *Annu. Rev. Pharmacol. Toxicol.* **49**, 243–263.
- Tan, M., Luo, H., Lee, S., Jin, F., Yang, J. S., Montellier, E., Buchou, T., Cheng, Z., Rousseaux, S., Rajagopal, N. et al. (2011). Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. *Cell* **146**, 1016–1028.
- Tanaka, S. (2001). Endocrine mechanisms controlling body-color polymorphism in locusts. *Arch. Insect Biochem. Physiol.* **47**, 139–149.
- Tanaka, S. (2006). Corazonin and locust phase polyphenism. *Appl. Entomol. Zool. (Jpn.)* **41**, 179–193.
- Tanaka, S. and Maeno, K. (2006). Phase-related body-color polyphenism in hatchlings of the desert locust, *Schistocerca gregaria*: re-examination of the maternal and crowding effects. *J. Insect Physiol.* **52**, 1054–1061.
- Tanaka, S. and Nishide, Y. (2012). Do desert locust hoppers develop gregarious characteristics by watching a video? *J. Insect Physiol.* **58**, 1060–1071.
- Tanaka, S. and Nishide, Y. (2013). Behavioral phase shift in nymphs of the desert locust, *Schistocerca gregaria*: special attention to attraction/avoidance behaviors and the role of serotonin. *J. Insect Physiol.* **59**, 101–112.
- Tanaka, S., Zhu, D. H., Hoste, B. and Breuer, M. (2002). The dark-color inducing neuropeptide, [His(7)]-corazonin, causes a shift in morphometric characteristics towards the gregarious phase in isolated-reared (solitary) *Locusta migratoria*. *J. Insect Physiol.* **48**, 1065–1074.
- Tariq, M., Wegrzyn, R., Anwar, S., Bukau, B. and Paro, R. (2013). Drosophila GAGA factor polyglutamine domains exhibit prion-like behavior. *BMC Genomics* **14**, 374.
- Umer, M. and Herceg, Z. (2013). Deciphering the epigenetic code: an overview of DNA methylation analysis methods. *Antioxid. Redox Signal.* **18**, 1972–1986.
- Uvarov, B. P. (1966). *Grasshoppers and Locusts*. Cambridge: Cambridge University Press.
- Uvarov, B. P. (1977). *Grasshoppers and Locusts*, Vol. 2. London: Centre for Overseas Pest Research.
- Vandegehuchte, M. B., Lemièrre, F., Vanhaecke, L., Vanden Berghe, W. and Janssen, C. R. (2010). Direct and transgenerational impact on *Daphnia magna* of chemicals with a known effect on DNA methylation. *Comp Biochem Physiol* **151C**, 278–285.
- Vandersmissen, T., Hoste, B., Baggerman, G., Huybrechts, J., De Loof, A., Chaltin, P., Proost, P. and Breuer, M. (2006). Degradation profile of [His(7)]-corazonin in the hemolymph of the desert locust *Schistocerca gregaria*. *Peptides* **27**, 539–548.
- Verlinden, H., Badisco, L., Marchal, E., Van Wielendaele, P. and Vanden Broeck, J. (2009). Endocrinology of reproduction and phase transition in locusts. *Gen. Comp. Endocrinol.* **162**, 79–92.
- Walsh, T. K., Brisson, J. A., Robertson, H. M., Gordon, K., Jaubert-Possamai, S., Tagu, D. and Edwards, O. R. (2010). A functional DNA methylation system in the pea aphid, *Acyrtosiphon pisum*. *Insect Mol. Biol.* **19** Suppl. 2, 215–228.
- Wang, X. and Kang, L. (2014). Molecular mechanisms of phase change in locusts. *Annu. Rev. Entomol.* **59**, 225–244.
- Wang, Y. and Leung, F. C. (2009). In silico prediction of two classes of honeybee genes with CpG deficiency or CpG enrichment and sorting according to gene ontology classes. *J. Mol. Evol.* **68**, 700–705.
- Wang, Y., Jorda, M., Jones, P. L., Maleszka, R., Ling, X., Robertson, H. M., Mizzen, C. A., Peinado, M. A. and Robinson, G. E. (2006). Functional CpG methylation system in a social insect. *Science* **314**, 645–647.
- Wang, H. S., Wang, X. H., Zhou, C. S., Huang, L. H., Zhang, S. F., Guo, W. and Kang, L. (2007). cDNA cloning of heat shock proteins and their expression in the two phases of the migratory locust. *Insect Mol. Biol.* **16**, 207–219.
- Wang, H., Yang, H., Shivalila, C. S., Dawlaty, M. M., Cheng, A. W., Zhang, F. and Jaenisch, R. (2013). One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* **153**, 910–918.
- Wang, X., Fang, X., Yang, P., Jiang, X., Jiang, F., Zhao, D., Li, B., Cui, F., Wei, J., Ma, C. et al. (2014). The locust genome provides insight into swarm formation and long-distance flight. *Nat. Commun.* **5**, 2957.
- Wei, Y., Chen, S., Yang, P., Ma, Z. and Kang, L. (2009). Characterization and comparative profiling of the small RNA transcriptomes in two phases of locust. *Genome Biol.* **10**, R6.
- Weiner, S. A., Galbraith, D. A., Adams, D. C., Valenzuela, N., Noll, F. B., Grozinger, C. M. and Toth, A. L. (2013). A survey of DNA methylation across social insect species, life stages, and castes reveals abundant and caste-associated methylation in a primitively social wasp. *Naturwissenschaften* **100**, 795–799.
- Whitman, D. and Ananthakrishnan, T. N. (2009). *Phenotypic Plasticity of Insects: Mechanisms and Consequences*. Enfield, NH: Science Publishers.
- Winston, M. L. (1987). *The Biology of the Honey Bee*. Cambridge: Harvard University Press.
- Wu, H. and Zhang, Y. (2014). Reversing DNA methylation: mechanisms, genomics, and biological functions. *Cell* **156**, 45–68.
- Wu, R., Wu, Z., Wang, X., Yang, P., Yu, D., Zhao, C., Xu, G. and Kang, L. (2012). Metabolomic analysis reveals that carnitines are key regulatory metabolites in phase transition of the locusts. *Proc. Natl. Acad. Sci. USA* **109**, 3259–3263.
- Wyatt, G. R. (1951). The purine and pyrimidine composition of deoxypentose nucleic acids. *Biochem. J.* **48**, 584–590.
- Wybrandt, G. B. and Andersen, S. O. (2001). Purification and sequence determination of a yellow protein from sexually mature males of the desert locust, *Schistocerca gregaria*. *Insect Biochem. Mol. Biol.* **31**, 1183–1189.
- Wynant, N., Verlinden, H., Breugelmans, B., Simonet, G. and Vanden Broeck, J. (2012). Tissue-dependence and sensitivity of the systemic RNA interference response in the desert locust, *Schistocerca gregaria*. *Insect Biochem. Mol. Biol.* **42**, 911–917.
- Xiang, H., Zhu, J., Chen, Q., Dai, F., Li, X., Li, M., Zhang, H., Zhang, G., Li, D., Dong, Y. et al. (2010). Single base-resolution methylome of the silkworm reveals a sparse epigenomic map. *Nat. Biotechnol.* **28**, 516–520.
- Xiang, H., Li, X., Dai, F., Xu, X., Tan, A., Chen, L., Zhang, G., Ding, Y., Li, Q., Lian, J. et al. (2013). Comparative methylomes between domesticated and wild silkworms implies possible epigenetic influences on silkworm domestication. *BMC Genomics* **14**, 646.
- Yamamoto-Kihara, M., Hata, T., Breuer, M. and Tanaka, S. (2004). Effect of [His(7)]-corazonin on the number of antennal sensilla in *Locusta migratoria*. *Physiol. Entomol.* **29**, 73–77.
- Yang, M., Wei, Y., Jiang, F., Wang, Y., Guo, X., He, J. and Kang, L. (2014). MicroRNA-133 inhibits behavioral aggregation by controlling dopamine synthesis in locusts. *PLoS Genet.* **10**, e1004206.
- Ye, Y. H., Woolfit, M., Huttley, G. A., Rancès, E., Caragata, E. P., Popovici, J., O'Neill, S. L. and McGraw, E. A. (2013). Infection with a virulent strain of *Wolbachia* disrupts genome wide-patterns of cytosine methylation in the mosquito *Aedes aegypti*. *PLoS ONE* **8**, e66482.
- Youngson, N. A. and Whitelaw, E. (2008). Transgenerational epigenetic effects. *Annu. Rev. Genomics Hum. Genet.* **9**, 233–257.
- Yu, M., Hon, G. C., Szulwach, K. E., Song, C. X., Zhang, L., Kim, A., Li, X., Dai, Q., Shen, Y., Park, B. et al. (2012). Base-resolution analysis of 5-hydroxymethylcytosine in the mammalian genome. *Cell* **149**, 1368–1380.
- Zemach, A., McDaniel, I. E., Silva, P. and Zilberman, D. (2010). Genome-wide evolutionary analysis of eukaryotic DNA methylation. *Science* **328**, 916–919.
- Zera, A. J. and Denno, R. F. (1997). Physiology and ecology of dispersal polymorphism in insects. *Annu. Rev. Entomol.* **42**, 207–230.
- Zhou, L., Cheng, X., Connolly, B. A., Dickman, M. J., Hurd, P. J. and Hornby, D. P. (2002). Zebularine: a novel DNA methylation inhibitor that forms a covalent complex with DNA methyltransferases. *J. Mol. Biol.* **321**, 591–599.
- Zwier, M. V., Verhulst, E. C., Zwahlen, R. D., Beukeboom, L. W. and van de Zande, L. (2012). DNA methylation plays a crucial role during early *Nasonia* development. *Insect Mol. Biol.* **21**, 129–138.







## 1.4

[illegible]

1.5

[illegible]

## 1.6

```
Trc Sirt4:      1  MSNLTRQLPFLKRTPILSDFIPRHSPAPSTDVQSLKNFIDDDYDKILVLTGAGISTESGIPDYRSEDVGLYARTNHRPVQYQEFKSEKVRQRYWARNYI 100
Lom Sirt4:      -----SAFVPEHNPNVNMEDFSKFMSFIETSRKIAVLTGAGVSTESGIPDYRSENVGLYARSSSRPVQFQDFLKSQVRRRYWARNFI

Trc Sirt4:     101  GWPKFSQTQPNITHYIVQFLESV-GKVGCVVTQNVDSLHFKAGSGNVIELHGTAFRVICLKCGKNYDRHKVQEKLIGLNPDLRETSQMIRPDGDVEISQE 200
Lom Sirt4:      GWPRFSSFSFNATHYALRDLEVIHHKIGFLVTQNVDDLHTKAGSNNIVELHGTAYNVICLNCDYNVSRQTFQEILQQINPNMKASTKMIRPDGDVELSQA
                ***.**      ** *** .. ** .  *. * .***** ** ***** *.*****. ***** * * *  ** * .**... ..*****.**

Trc Sirt4:     201  KVENFKPPFCDHCQGVLKPDITFFGDNVPKERVDKVRQNVQTQSDAILVLGSSLSVFSGYRIILQAVEEEKKDVAIVNIGPTRADKHAKLKISARCGDILPR 300
Lom Sirt4:      EAETFGVPECPCTCGGILKPDIIFFGENVPRMRVETVKRIVEDADSLVLGSSLSVFSGFRIVLQAAELEKPIGIINIGKTRGDSYAHIKVNGKCGDIVPR
                . * * * * * *.***** ***.***. **. *.. *  ..*****.***.*** * . * . * .*** ** * . * .*. .****.**

Trc Sirt4:     301  IFE 302
Lom Sirt4:      ILD
                * .
```

**Fig. S2: Putative sequences of Histone Acetyltransferases found in the genome of *Locusta migratoria*. Sequences were derived from concatenated BLAST hits.**

## 2.1

```

Drm Mof:      1  MSEAELEQTPSAGHVQEPIEEEHEPEQEPTDAYTIGGPPRTPVEDAAAEELSASLDVSGSDQSAEQSLDLSGVQAEAAAEESEPPAKRQHRDISPISEDST 100
Drm Mof:     101 PASSTSTSSSTRSSSSSRYYDDVSEAEAEAPPEPEPEQPQQQQQEEKKEDGQDQVKSPGPVELEAQEPAQPQKQKEVVDQEIETEDEPSSDTVICVADINPYG 200
Drm Mof:     201 SGSNIDDFVMDPDAPPNAIITEVVTIPAPLHLKGTQQLGLPLAAPPPPPPPAAEQVPETPASPTDDGEEPPAVYLSPIYRSRYMQESTPGLPTRLAPRD 300
Drm Mof:     301 PRQRNMPPPAVVLPITQTVLSANVEAISDDSETSSSSDDDEEEEDDEDDALTMEDHNTSRETIVITGDPMLQKIDISENPKIYFIRREDGTVHRGQVLQS 400
Drm Mof:     401 RTTENAAAPDEYYVHVGLNRRLDGVVGRHRISDNADDLGGITVLPAPPLAPDQPSTREMLAQAAAAAASSERQKRAANKDYLLSYCENSRYDYSR 500
Drm Mof:     501 KMTRYQKRRYDEINHVQKSHAELTATQAALKEHESESITKIYIDKLQFGNYEIDTWYFSPFEEYGKARTLYVCEYCLKYMFRSSYAYHLHECDRRRPP 600
Lom Mof:      -----IEFGQYEIQTWYSSPYPQYARCIISLFLCEFLKYTKSKAVLERHQDKCTWRHPP
                        . . . * * * * * . . . * * * * * . . . * * * * *
Drm Mof:     601 GREIYRKGNISIIYEVNGKEESLYCQLLCLMAKLFLDHKVLYFMDPFLFYILCETDKEGSHIVGYFSKEKKSLENYNVACILVLPHPQRKGFGKLLIAFS 700
Lom Mof:      ATEIYRHKDLSVFEVDGNVNIYCQNLCLLAKLFLDHKTLYDVEPFLFYVLTKNDDK-----
                        * * * * . . . * * * . * * * * * * * * * * * * * * * * *
Drm Mof:     701 YELSRKEGVIGSPEKPLSDLGRLSYRSYWAYTLLLELMKTRCAPEQITIKELSEMSGITHDDIIYTLQSMKMIKYW-KGQNVICVTSKTIQDHLQLPQFKQ 800
Lom Mof:      YLLSKQEGQPGTPEKPLSDLGRVSYHAYWKSIVLEYLDQH-RNEPFSVDAVCQETGVYSHDVAATLQMLGMVRRCPDGRLLAIDWAQVDAHASRVRASR
                        * * * . . . * * * * * * * * * * * * . . . * * * * * * * * * * * *
Drm Mof:     801 PKLTIDTDYLVWSPQTAAAVVRAPGNSG 827
Lom Mof:      TRIPLDPECLRWTPLVATVPTR-----
                        . . . * * * * * * *

```

## 2.2

```

Drm Gcn5:      1  MSGGPSITIKSQPIDGNNTGNAAAQQQQQAANGAATAGASGAAGSAQNPGHGGGAASGASVPAEGTRQNSLQRIQQRKQKVFNLVPVPQKLAKLSMYSACQ 100
Drm Gcn5:     101 SEGCRCTGWKTPQENRHRDVESSYCPEFNEECRNTSCRHSLRSHIAHLDN-ISSSSMNELLGAIIDMENLFMSMQRVEDEDTKKVYQYLFRLLRQCVLTR 200
Lom Gcn5:      -----THVSHLQPLSMEEVNQLLGMVVDVENIFMSMHREEDADTKRVYVYLFQLLRKCILTM
                        . . . * * * . . . * * * * * * * * * * * * * * * * * *
Drm Gcn5:     201 QQAVIRGPLGDPFFETPCITKAVLSLVFYKYNHLSTPELQTMTEVAKTFLNFLNHYNFESPSTRGDLTHEDASNYKINYTRWLVFCHVPAFCNSLRQCE 300
Lom Gcn5:      TKPSVEGPLGQPPFERPSIAKAITNFVLYKFGHL--EWQTMYDLAKMFLHCLNHNWFETPSVRRHTVTAEEISAYKVNYTRWLVFCHVPAFCDSLPHFE
                        . . * * * * * * * * * * . . * * * * * * * * * * * * * * * * * * * * * *
Drm Gcn5:     301 TSLVFGRTLRLTVFQCMSQQLKKKCIISERDRFPEDKRSIITLMPKFLETTLRAELLKDDSPIWDTSYRPSNSFVIQQRKRNOEVANVPIGPSAASIGGNKR 400
Lom Gcn5:      TTLVFGRTLRLRAVFRSVCRQLMDKCHSERDRMPPEKRVLV-----
                        * . * * * * * * * * * * * * * * * * * * * *
Drm Gcn5:     401 TSVGEPLHKRIKKEPTDRPSSSENDDLPADVVMRAMKSVSESKTTNKAEILFPVNVSRDENVKAEQKRAIEFHVVGNSLTKPVDKQTVLWLLGLQLVFA 500
Lom Gcn5:      -----RTRLPPQAVFPENAPRDETAQEEARGVISFHVVGNSLMRPVSKSTMVLWLVGLQNVFS
                        . * . . * * * * * * * * * * * * * * * * * * * * * *
Drm Gcn5:     501 YQLPEMPREYISQLVFDTKHKTLALIKENQPIGGICFRFPFSQGFTIEIVCAVTMSEQVKGYGTHLMNHLKDYSIQRGIKHLLTFADCDAGYFKKQGFS 600
Lom Gcn5:      HQLPRMPKEYISRVLFD-----GYFKKQGFS
                        . * * * * * * * * * * * * * * * * * * * * *

```





**Fig. S3: Putative sequences of DNA methyltransferases found in the genome of *Locusta migratoria*. Sequences were derived from concatenated BLAST hits.**

### 3.1

```

Apm Dnmt1a:      1 MKSNTSKFKQTSIIDMFSQQLAQKKQMNDATNKEDIQKLQNIKTENNILQKCEQNKENHDPKTDEEYLDKHIEKKAKLTDVNILPTPQIKNINIFHNKC 100
Lom Dnmt1:      -----KRAKVLVDYKSQQLAKEQLIREEETLQRQRSRSPHTRNTGSDNSSCENSEEEVEDEEKELEDEEEKE-----VRSINL--EKC
                  * . . * * * * * . . . * . . . . . * * . * . * * . * * . . . * * . * *
Apm Dnmt1a:    101 EYCRQKLN--EIKFYPGHPNGAVDEEIALIDPKLCLFNGTESFIHESDERPQNKLTYFSVYDKNGHLCSDFTGLIEKNVMLYFSGYIKAIYEEDASPEGG 200
Lom Dnmt1:      KTCHQLLSNPCLKLYEGHPEGAVEEFVALTDPKLSLFSGQEETVHEHDERPQNKITHFSIYDKEGHICPFDGGIERNGLLYFSGYMKAVYEENPDPEGG
                  . * * * . * . * * * * * * . * * * * * * * * . * * * * * * * * . * * * * * * * * . * * * * * * * *
Apm Dnmt1a:    201 VPTKDMGPINNEWVSGFDGGELALIGFNTAFAYIILMDPSEYAPFMDIVKEKIYMSKLVIEFLLDEINPSYEDLLNKFQTIVPPKNMSKFTEDSLLRHA 300
Lom Dnmt1:      IPTKDMGPINNEWWISGFDGGELALIGFSTAYGDYYLMEPNESYSPYMDAVRQKIFLSKAVIEFLLTEIDPSYEDLLNKLQTVVPPSGISCFSEDILLQHA
                  . * * * * * * * . * * * * * * * * . * * * * * * * * . * * * * * * * * . * * * * * * * *
Apm Dnmt1a:    301 QFICDQVLSFDN-SADSEDTLLITSPCMRALANLAGVTLGKRVTFRRTQREQ--KVRKPAWTKATTTKLVSDFENIFTDQLIKHDDKITMGPKRQRCGV 400
Lom Dnmt1:      QFVCDQVMSFDSASEDTDDQLLIATPCMRSLIKLAGVTLGKQAMRRAERDVTKPKKSEWTKATTTNLVHEVFETFFTEQIDRDNNK--GPKRRRCGV
                  * * . * * * . * * . * * * . * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
Apm Dnmt1a:    401 CEHCQQPDCGLCNACKDMIKFGGTGRSKQACVKKRCPNMAIQEADSDPENEDQYEILMENKKVE---EEKSKKKDFKELKKDIIWIGDQIANDNLKTTY 500
Lom Dnmt1:      CEACQSPDCGTCIACKDMVKFGGSGRSKQACVQRRCANMAVEAAEESDVEDDNDLNVKVKSSCLKIGLSENQKKIVSRDKRSVLWIGDMITKVGPFKY
                  * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
Apm Dnmt1a:    501 KSVIVGDEEIGINDYVLVEPRNPAIPSHVAKVIYMWENKNGIKQFHANWLHRGNDTILGETSDPIELFLSDDCDDIPFKSVRSKCTVIFKEIQKNWNELG 600
Lom Dnmt1:      RAFSINDEEFKVNDYVIVNPLPATPVLIAKISYMWEDKGSKWFWHATW--RGTDTVLGETADPLELFVDECENTLISTVKCKANVHFTPIPSNWRREL
                  . . . * * * . * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
Apm Dnmt1a:    601 NMDLNFENDIKDLGDKTFFFQKRYIPETARFEDPVPDVLYLHKKNSYRFCPACTRLNMLEQFNTPKVYERIEEINSKEVIYGLVKYKGEDYRVGTTVFLQ 700
Lom Dnmt1:      GATTNKENNFED-DEKTFYQKR-----
                  * * * . * * * * * * * *
Apm Dnmt1a:    701 PDVFNFKYKTTYQDVLKLLKENVDEDMYFEFYRKSSDHIKGSNFDTPDPFCIGYINKIFANTNDMLVSPSDINIMVNKLYRPENTHKSSTLMEQADLNMV 800
Lom Dnmt1:      -----VDEDLYPEYRKTSDFVKGSEETPEPFSIGYISSVFIHSSDIIA--RDIKLRVHKLYRPENTMKDREAAQADLNKL
                  * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
Apm Dnmt1a:    801 YWSNEVCNIKFIKVVGKCYLTYSENLNQSIIEEWIAAGPYRFYFNEAYNVQEKTFDEPPYHAISIGKCGSKGNLKFKGKTEESEKRAIVNTPIEYKKIS 900
Lom Dnmt1:      YWSDEGCVISFDNVEGKCYVYSENLDVTVEWTKGGPDRFYFSEAYNSAEGTFDEPCHKATLIGSQGKGKGGKSSDETRK--VENKYKDWPKIS
                  * * * . * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * * *
Apm Dnmt1a:    901 EKLKTLDFVAGCGGLSEGLHQAGVAENLWAEKEESAAYARLNNPNATVFIEDCNVLLKKVMNGETTNEIGQKLPQKGQVELLCGGPPCQGFSGMNRFN 1000
Lom Dnmt1:      SKLKSLDVFAGCGGM-----
                  * * * . * * * * * * *
Apm Dnmt1a:    1001 SRQYSLFKNSLVVSYLSYCDYRPNFFIMENVNRNFVSFKRSMVLKLTLRCLIRMGYQCTFGILQAGNYGIPQTRRRMIILAAAPGQMLPKYPEPSHVFSK 1100
Lom Dnmt1:      -----KCTISLWPRSSFSLLFSRA--IILAAAPGEVLNPFPEPTHVFS
                  . * * . . . . * * * * * * * * * * * *

```







Fig. S5: Putative sequences of Histone Methyltransferases found in the genome of *Locusta migratoria*. Sequences were derived from concatenated BLAST hits.

## 5.1

```

Drm Set:      1  MSSVPKRAKLDGAPADGNTSAAAGNNEEESEALEQIDACQNEIDALNEKASEEILKVEQKYNKLRKPCYEKRSELVKRIPNFWVTSFINHPQVSGILDEE 100
Lom Set:      -----KQRKSEGSAAE--TEDERDYDNETQKALEEIDACQNEIDALNEKASEEILKVEQKYNKLRKPYFEKRNEIIKRIPNFWVTAFAVNHPQIAAILEEE
                * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . * . *
Drm Set:     101  EEECLHALNKLEVEEFEDIKSGYRINFHFEDENPYFENKVLTKFEHLNSAAAASENGDWPASTSTPIKWKEGKNLLKLLLTTPYGNKKKRNSEYKTFDFWFS 200
Lom Set:     EEDCLHYLTKEVEEFEDIKSGYRINFHFENPYFENNILTKEFHLGTTGKLS-GD-PASQSTQIRWKDGHDLTKKAKEKTAAGKRKRDLDQKTFHFHWF
                ** . *** * ***** . ***** . ***** . .          ** *** ** * . ** . * *          *          . ** . . **** **
Drm Set:     201  DNTDPVNDEIAELIKDDLWPNPLQYYLVPDIEVEPEDEEDNEDNDEEAFDDEDGEDGEDGEGEEEEDEDDK 269
Lom Set:     DHGDPAADDIAEV-----
                * . **      * . ***

```

## 5.2

```

Peh MLL:      1 MEIVSEDESEDLSQPPKTPDIHLSDTEDVPTPSRLKSKKSSAMEALAALGLDEFVGEIKIKERKENKKEKLSGFAGIDRKLDSDSRSDVNTTEKKSQNK 100
Peh MLL:     101 NADKNKYKDVTVNDHKSEKFNDRTNELNDKRKRQKMEIINGDHYSEKVSDDLVDNVSKYGSVKLGKDEALEIDVEDEPSRSSPESQIQMEHSYSLPRER 200
Peh MLL:     201 DPSSSNSPASESEIKKKQEAFTLDHDYTTKSKSPLTPKLDDNKFGKTTVKEIKKSSEKLISHLLDTYLPKEKVTVPVKYKERNLIEEMTVLYEFLTKGIDA 300
Peh MLL:     301 EDIGYLKRSYDSMLADESQAYWLNDTHWVDHPVTDLSSGVPSKKARVHSTGSARTQGYKIDAVEKMRHKHHYKNTVQNQNGGTAHGIRGVAELVKGVNA 400
Peh MLL:     401 ASREARSNQRRLLTAFGTATDSDLKFNQLKFRKKQLKFSKSDIHDWGLFAMEPIAADEMVIEWVGQVMRPFPLADFREKEYEYKRGIGSSYLFRIDVDTII 500
Lom MLL:      -----QFRKKQLKFAKSAIHDWGLFAMEPIAADEMVIEWVGQVMRPLVADLRERHYEATGIGSSYLFRIDVDTII
                .*****. ** *****. ** .**.*. ** *****.***

Peh MLL:     501 DATKCGNLARFINHSCNPNCYAKIITIEGQKKIVIYSKKDIKVDEEITYDYKFPIEEEKIPCLCGAAQCKGYLN 574
Lom MLL:      DATKCGNLARFINHSCNPNCISRIIPGSSRRHIAIYSKRDIQVGEEVTDYDYKFPIEPEK-----
                ***** ** * * * * * * * * * * * * * * * *

```

### 5.3

Peh	E (z):	1	MNSTKVPPEWKRRVKSEYIKIRQQKRYKRADEIKEAWIRNWDENHNHVQDLYCESKVWQAKPYDPHPVDCVKRAEVTSYNGIPSGPKQVPICVINAVTPI	100
Peh	E (z):	101	PTMYTWAPTQQNFMVEDETVLHNIPYMGDEVLDKDGKFI EELIKNYDGKVHGDKDPSFMDDAIFVELVHALMRSYSKELEEAAPGTATAIKTETLAKSKQ	200
Peh	E (z):	201	GEDDGVDVDADGESPMKLEKTDSDKGLTEVEKKETEEPLETEDADV KPDVEEVKDKLPFPAPIIFQAI SANFPDKGTAQELKEKYIELTEHQDPERPQE	300
Peh	E (z):	301	CTPNIDIGIKAESVSRERTMHSFHTLFCRRCFKYDCFLHRLQGHAGPNLQKRRYP ELPKFAEP CSNSCYMLIDGMKEKLAADSKTPPIDSCNEASSED SND	400
Peh	E (z):	401	SNSQFSNKDFNHENSKDNGLTVNSAAVAEINSIMAGMMNITSTQC VWTGADQALYRVLHKVYLKNYCAIAHNMLTKTCRQVYEFAQKEDAEFSFEDLRQD	500
Lom	E (z):		-----TVSQPTSTRTSFSLGLMGPEET-AEW TGADQSLFRALHKVFLNNYCAIAQIMLTCTCQVYEFAQKESADIITEESLKD ** . * . * . * . * . * . * . * . * . * . * . * . * . * . *	
Peh	E (z):	501	FTPPRKKKKKQRLWSLHCRKIQLKKDSSSNHVYNYTPCDHPGHPCDMNCSCIQTQNFCEKFCNCSSDCQNRFP GCRCKAQCNTKQPCPYLAVRECDPDL C	600
Lom	E (z):		FTPPRKKKKKHLWSMHCRKIQLKKDSSSNHVYNFTPCDHPNPQCDQLCPCIG AQNFCEKFCQCSSDCQNRFP GCRCKAQCNTKQPCPYLAVRECDPDL C ***** . *	
Peh	E (z):	601	QACGADQFKLTKITCKNVCVQRGLHKHLLMAPSDIAGWIGIFLKEGAQKNEFISEYCGEII SQDEADRRGKVYDKYMC SFLFNLNND FVVDATRKGNKIRF	700
Lom	E (z):		VTCGAEQFDITKISCKNVSVQRGLRK----- *** * . * . * . * . * . * . *	

Peh E(z): 701 ANHSINPCYAKVMMVTGDHRIGIFAKRAIQPGEELFFDYRYGPTEQLKFVGIEREMEIV 760  
Lom E(z): -----

## 5.4

[illegible]

## 5.5

[illegible]

**Fig. S6: Putative sequences of Histone Demethylases found in the genome of *Locusta migratoria*. Sequences were derived from concatenated BLAST hits.**

## 6.1

```

Drm Su(var) 3-3: 1 MKPTQFGGSSSKMTEPIEYVTLISDDSDGEPTPKRNVNHPPSALSAPNPGQKQKHPDEDSNDAPATSDERRTSRRNRPKVDYSNRPSGSGDTASNDKSGS 100
Drm Su(var) 3-3: 101 ASMGPNNQAEERRSQSQRKSEANATSSSVSGPSAGNSRPSQNGDSKDRDAGTPTVLSGQEGAVFQSRLPFNKMTPNEEACFPDISRSGILGHRVFLNIR 200
Lom Su(var) 3-3: -----AGLEGAAFQSRLPFDKLTANEASCFPDVAQGGPPQTQKVFLHIR
                                     . * * * * . * * * * * . * * * * . * * * * *
Drm Su(var) 3-3: 201 NSLLHMMVDNPKIQLSFEIALKNLPPFPDSEPSLVRVRHVSFLERHGFINGIFKRLKPIPAKKLGKVVIVIGAGISGLAVAHQLQQFGM-DVIVLEARDRV 300
Lom Su(var) 3-3: NRLLQLWLENPKQQLLLENAL-----INFFFFHFRHRT-----MIGVLGGGISGLAAHYLLKTKVQPVIIILEASHRI
      * * * * * . * * * * * . * * * * * . * * * * * . * * * * *
Drm Su(var) 3-3: 301 GGRISTFRKNSYIADVGAMVVTGVYGNPMTILSKQIGMDLVPIQQTCPLYPGDPGKVPVPEKDDVIREFNRLLESASYLSHRLDFNYAGDCPVSLGDALE 400
Lom Su(var) 3-3: GGWIQTTSNPT-----GN-----
      * * * * * . * * * * *
Drm Su(var) 3-3: 401 WIISMQEMQVMHKGQHMQEIIATQTKIIEQRRRLKTLRDTIGKLKNEHLAMINQRKPKGTDGDLKYCYQEFNIRNTQIKMEETISTFHDHAEKQMLA 500
Lom Su(var) 3-3: -----IYEHGPRTIRVRGPLG---ENTLLL-----VKELGLSNK-----VLPIRS
                                     * * * * * . * * * * * . * * * * *
Drm Su(var) 3-3: 501 KLHELEQNRPSDVYLSSRDRLILDWHFANLEFANATRLNNLSLKHWQDDDDFEFIGHHTTVRNGYSCVPVALTENLDIRVNSAVKEIKYGTGKGEVVAEN 600
Lom Su(var) 3-3: THPELDASPPSDVYLSPDRQILDWHFANLEFANATPLDKLSLKHWQDDDDFEFTGSHLTVRNGYSCVPVALSEGLDIKLTAVRHIRYTTSGVEV----
      . * * * * . * * * * * . * * * * * . * * * * * . * * * * * . * * * * *
Drm Su(var) 3-3: 601 LKTSNSQMTYKADLVVCTLTLLGVLKVAVAHKESQQSNTVKFDPPLPDWKQQAIRKLGFGNLNKVVLCFDRIFWDPNANLFGHVGSTTASRGEMFLFWSIS 700
Lom Su(var) 3-3: -----SYADHIICTVPLGVK-----ERASTM-FCPALPQYKTESINRLLFGTVQVVLCFDRIFWDPSTHLFGHVGSTTASRGELFLFWNLY
      . * * * * . * * * * * . * * * * * . * * * * * . * * * * * . * * * * *
Drm Su(var) 3-3: 701 SSPVLLALVAGMAANLVESVTDIIIGRCMSVLKNIFGNTSVPQPKETVTVTRWRSDPWARGSYSYVSVGSSGSDYDLLAAPVIPSSSKDAEGLPRLFFAG 800
Lom Su(var) 3-3: RAPVLLALVAGEAAAIMENVSDDIVGRCIAVLKGIFGNNSVPQPKETVTVTRWRADPWSRGYSYFVAVGSSGADYDILAAPVTP-----PRLFFAG
      . * * * * * . * * * * * . * * * * * . * * * * * . * * * * * . * * * * *
Drm Su(var) 3-3: 801 EHTIRNYPATVHGAYLSGLREAGRIADYYLGYPEGTPPDIGYSVAEAAANLVSVGNVVKLRDLSPNLSDSPSSSKKSEENSNSNTADSTELQ 890
Lom Su(var) 3-3: EHTIRNYPATVHGAFLSGLREGGRIADYFTGCPYA-PP-----
      * * * * * . * * * * * . * * * * * . * * * * *

```

## 6.2

```

Apm Su(var) 3-9: 1 MGGEELGVTTGQPNLYKQDLSKLDVSKLTALSREVISRQATINIGTIGHVAHGKSTIVKAISGVQTVRFKNELERNITIKLDTAEDSTRGEQEANLG 100
Lom Su(var) 3-9: -----GTIGHVAHGKSTVVKAIISGVQTVRFKNELERNITIKLTENTTLGI-----
                                     * * * * * . * * * * * . * * * * *
Apm Su(var) 3-9: 101 TNKMSTRGKRPPASPGYVQPPTKHQKLDQKFIFPQENNYNDNYF-----YKSNGSNSSNSDVLFKQNKEEEQTINRKNSDYLDNQEVSMENKSC- 200

```

Lom Su(var) 3-9: -----ETLFPCENNYSNQGTCINTNSQYKNPEASLNDNSNTELKHNL---AHIELDNESFVNERN-EVSDQTNSECQ  
. . \*\* \* . . . . . \* . . . . \* . . . . . \* . . . . . \* . . . . . \* . . . . .

Apm Su(var) 3-9: 201 -----TDSIEKYKMFENLVKLKELRIILEDIKHINTRHNTKNGMKTLLSETDIWEVEQILAKKEIKGVPTYLIKWNWDLKYNTWEPIS 300  
Lom Su(var) 3-9: NDYKIIIEVSNKNFNDSIQTEVDIFVDINNEEF-----NKLSEVLDRKHVEVEKVIDHCTSKNKRYYYVKWKGWSDCYNTWEPEE  
. \*\* . . . \* . . . . . \* . . . . . \* . . . . . \* . . . . . \* . . . . .

Apm Su(var) 3-9: 301 NLINCSDILEEFERNRLQLLESFKRKVNFPNNQDI-----EKFL-NYLKRGGKTLTSSIVESNTMFINILKFLKQ---KYVKNKLEKVI 400  
Lom Su(var) 3-9: HLKTCTSVIKTY----IQFLDRINSGNLYSTQEAKDLQEKLVSVSKEYELEKLMKLYVKSSGIPYTVCSSEE--VVHKMKMALRPSKKETNGEKNILKV  
. \* . \* . . . . . . \* . \* . . . \* . \* . . . \* . \* . . . \* . \* . . . \*

Apm Su(var) 3-9: 401 KHDILRMLIIDLRKKQLKSLEDWENEMNIITKGKPLIRVENIIDLEGAPQNFYIEEYLPNGVVIIPDDPPIGCECKTC-N-SKTKCCFAQDDGLCPYTL 500  
Lom Su(var) 3-9: KNDILVRIYDIYRMVQLDALKTWEKELNDKATEKAKIFVENNVLDLEGGPSNFTYINDYIPGKGVVIIPDDPPIGCACGPKCDRHSKECCGQSGSIFAYSH  
\* . \*\*\* . \* \*\* . \*

Apm Su(var) 3-9: 501 KHKIRVPPGTPPIYECNKRCNCDIDCINRVVQRGTMQFCIFRTANGRGWGVKTMKTIKKGFSFVTQYVGEVITNEEAERKRGKEYDAAGRTYLFDLDYNESE 600  
Lom Su(var) 3-9: KHLVADVGVPIYECNKRCCTQDCPNRVVQKGRKIDLSIFRTSTGCGWGVKTLQYIKKGSFICEYVGEVIETKEAESRGKVYDAEGRTYLFDLDFN--V  
\*\* . . \*

Apm Su(var) 3-9: 601 EQCPYTVDAAIYGNISHFINHSCDPNLAVYGVWINCLDPNLPLKALFATKDIKQNEEITFDYMCQSSKNS--ENSIMQRASMKENLVNYPEFQENVQLCS 700  
Lom Su(var) 3-9: DACPYTVDAAYGNVSHFINHSCDPNVAVYAVWVNCLDPNLPLKALFAKNDINVGDELTFDYMHNNLSIQDDVNSIANR-----VVSPPRSRLLLPN  
. \* . \*

Apm Su(var) 3-9: 701 EISESYSSNNKTLCKCDAQNCRRYLF 683  
Lom Su(var) 3-9: ADNSTDRTFYKTPCKCGAKQCKRYLF  
. . . . \* . \* . \* . \* . \* . \*



**Table S1:** Examples for differences in morphology, physiology, behaviour, and molecular changes between solitary and gregarious locusts of *Schistocerca gregaria* FORSSKÅL 1775 and *Locusta migratoria* (LINNAEUS 1758).

<i>Schistocerca gregaria</i>		<i>Locusta migratoria</i>	
Solitary	Gregarious	Solitary	Gregarious
<b>1. Morphological differences</b>			
Large females, small males	Smaller females, large males	Large females, small males	Smaller females, large males
Long wings	Short wings	Long wings	Short wings
Convex pronotum	More concave pronotum	Convex pronotum	More concave pronotum
High F/C ratio <sup>1</sup>	Low F/C ratio	High F/C ratio	Low F/C ratio
Low E/F ratio <sup>2</sup>	High E/F ratio	Low E/F ratio	High E/F ratio
More sensilla on antennae	Less sensilla on antennae	More sensilla on antennae	Less sensilla on antennae
More ovarioles	Less ovarioles	More ovarioles	Less ovarioles
More, smaller eggs	Fewer, bigger eggs	More, smaller eggs	Fewer, bigger eggs
Light hatchlings	Dark hatchlings	Light hatchlings	Dark hatchlings
<i>Beige adults</i>	<i>Yellow adults</i>	<i>Green adults</i>	<i>Darkbrown females/ yellow males</i>
<i>Green/beige nymphs</i>	<i>Black and yellow nymphs</i>	<i>Green/lightbrown nymphs</i>	<i>Black and orange nymphs</i>
<i>5-6 nymphal stages</i>	<i>5 nymphal stages</i>	<i>5-7 nymphal stages</i>	<i>5 nymphal stages</i>
Long hind legs	Shorter hind legs		

5-6 eye stripes	5 eye stripes
Large eyes	Small eyes
Small brain	Larger brain (30%)
Long antennae	Short antennae
More sensilla on hind legs	Less sensilla on hind legs
More frontal hairs	Less frontal hairs
Less touch receptors	More touch receptors

Shorter sternal hairs  
Less spermatophores

Longer sternal hairs  
More spermatophores

---

## 2. Physiological differences

---

Long-lived	Short-lived	Long-lived	Short-lived
Long maturation times	Short maturation times	Short maturation times	Long maturation times
Lower O <sub>2</sub> consumption	Higher O <sub>2</sub> consumption	Lower O <sub>2</sub> consumption	Higher O <sub>2</sub> consumption
Lower heartbeat frequency	Higher heartbeat frequency	Lower heartbeat frequency	Higher heartbeat frequency
Lower fat content	Higher fat content	Lower fat content	Higher fat content
Lower carbohydrate content	Higher carbohydrate	Lower carbohydrate content	Higher carbohydrate
Lower spontaneous activity of TCD in the light <sup>3</sup>	Higher spontaneous activity of TCD in the light		
No difference between activity of TCG in light and dark <sup>4</sup>	Higher spontaneous activity of TCG in the dark		

### 3. Molecular differences

<i>Lower SGPP1-4 expression<sup>5</sup></i>	<i>Higher SGPP1-4 expression</i>	<i>Higher LMPP2 expression<sup>6</sup></i>	<i>Lower LMPP2 expression</i>
<i>No difference in octopamine levels</i>	<i>No difference in octopamine levels</i>	<i>Higher octopamine levels</i>	<i>Lower octopamine levels</i>
<i>Lower JH titer<sup>7</sup> (debated)</i>	<i>Higher JH titer (debated)</i>	<i>Higher overall JH biosynthesis</i>	<i>Lower overall JH biosynthesis</i>
Low ecdysteroid level in eggs	High ecdysteroid level in eggs		
High ecdysteroid levels in haemolymph	Low ecdysteroid levels in haemolymph		
No phenylacetonitrile production	Phenylacetonitrile production		
Low yellow protein (YP) levels	High YP levels in adult males		
Low carotenoid/biliverdin ratio	High carotenoid/biliverdin ratio		
Low phase-related peptide (PRP) level	High PRP level		
Low levels of PRP in eggs	High levels of PRP in eggs		
Slower corazonin breakdown	Faster corazonin breakdown		
High expression of NPP1-2 <sup>8</sup>	Low expression of NPP1-2		
Low expression of NPP3-4 <sup>8</sup>	High expression of NPP3-4		
Higher levels of 8 neurochemicals: Arg, Asp, Glu, Gly, taurine, GABA, dopamine, and serotonin	Higher levels of 4 neurochemicals: acetylcholine, citrulline, N-acetyldopamine, and tyramine		



Higher levels of 155 metabolites, especially phosphatidylethanolamines (PEs) and diacylglycerols (DGs)	Higher levels of 164 metabolites, especially carnitines, lysophosphatidylcholines, several carbohydrates, amino acids, and some DGs		
Differential expression in 214 genes (long-term solitary), 114 genes upregulated, including genes involved in energy metabolism, protein synthesis, and antioxidant genes	Differential expression in 214 genes (long-term gregarious), 100 genes upregulated, including heat-shock and immunity genes	Upregulation of genes involved in energy metabolism and protein synthesis	Upregulation of 6 heat-shock protein (HSP) genes
		Lower levels of Neuroparsin A	Higher levels of Neuroparsin A
		Lower OMP levels in mature adults <sup>9</sup>	Higher OMP levels in mature adults
		Higher levels of APRP <sup>10</sup>	Lower levels of APRP <sup>10</sup>
		Lower levels of dopamine	Higher levels of dopamine
		Higher haemolymph levels of lipid, trehalose, and putrescine	Lower haemolymph levels of lipid, trehalose, and putrescine
		Differential expression in 532 genes (2-3 generations), including upregulation of mostly anabolic, biosynthetic, and muscle-specific gene expression	Differential expression in 532 genes (2-3 generations), including very strong upregulation of the JHPH gene family <sup>11</sup> in the head
			Upregulation of a cluster of synaptic transport components, neurotransmitter and neuro-modulator receptors, neurotransmitter synthetases, and G protein coupled receptors

Upregulation of *take out* in antennae at 4<sup>th</sup> instar

Upregulation of genes for CSPs (chemosensory protein) at 4<sup>th</sup> instar

Upregulation of genes of general metabolism and catecholamine biosynthesis at 4<sup>th</sup> instar

Differential expression in 4893 genes in the brain (short-term, 64h), including upregulation of genes in redox biology

Differential expression in 4893 genes in the brain (short-term, 64h), including upregulation of genes in synaptic transmission, carbohydrate metabolism and nucleosome assembly

Higher abundance of longer (>26-29 nt<sup>12</sup>) small RNAs

Lower abundance of longer (>26-29 nt) small RNAs

Higher expression levels of micro-RNA 133

Lower expression levels of micro-RNA 133

Upregulation of transposon element I in antenna and labial palp at 4<sup>th</sup> instar

Upregulation of transposon element I in brain and wings at 4<sup>th</sup> instar

45 alternative spliced isoforms between gregarious and solitarious animals

90 genes differentially methylated

---

#### 4. Behavioural differences

---

Flight at night

Flight by day

Flight at night

Flight by day

Slow, creeping gait	Rapid, upright gait	Slow, creeping gait	Rapid, upright gait
Restricted diet	Generalists	Restricted diet	Generalists
Avoid other locusts	Attracted to locusts	Avoid other locusts	Attracted to locusts
Relatively sedentary	More active	Relatively sedentary	More active
Less intense mating behaviour	More intense mating behaviour	Less intense mating behaviour	More intense mating behaviour
Lower grooming frequency	Higher grooming frequency	Lower grooming frequency	Higher grooming frequency

---

Similar findings for both species are listed first, followed by differences (*italic*) and findings that have only be described in one of the species. Abbreviations: <sup>1</sup> F/C (length of femur/ maximum head width), <sup>2</sup> E/F (length of fore-wing/length of femur), <sup>3</sup> TCD (tritocerebral dwarf), <sup>4</sup> TCG (tritocerebral giant), <sup>5</sup> SGPP (Schistocerca gregaria pacifastin-related precursor), <sup>6</sup> LMPP (Locusta migratoria pacifastin-related precursor), <sup>7</sup> JH (juvenile hormone), <sup>8</sup> NPP (neuroparsin precursor), <sup>9</sup> OMP (ovary maturing parsin), <sup>10</sup> APRP (adipokinetic hormone (AKH) precursor-related peptide), <sup>11</sup> JHPH gene family (juvenile hormone binding protein, hexamersins, prophenoloxidase, and hemocyanins), <sup>12</sup> nt (nucleotides)



**Table S2: List of genes found.** Evidence of the existence of genes with putative roles in epigenetics in the genome of *Locusta migratoria*. Sequences of known genes were used to search the recently released genome sequence. Hits with significant similarity were then reblast to the genome database of NCBI. When this resulted in hits of the same (orthologous) gene, the hit in the *Locusta* genome was assumed to be the orthologue of the initial query sequence.

Group	Gene	Fig reference
1. Histone deacetylases (HDAC):	<i>Rpd3</i>	1.1
	<i>HDAC3-like</i>	1.2
	<i>HDAC5-like</i>	1.3
	<i>HDAC6-like</i>	1.4
	<i>HDAC-X</i>	1.5
	<i>Sirt4</i>	1.6
2. Histone acetyltransferases (HAT):	<i>Mof</i>	2.1
	<i>Gcn5/Pcaf/Kat2A</i>	2.2
	<i>Taf1</i> : not found	
	<i>NCOA2</i> : not found	
3. DNA methyltransferases (Dnmt):	<i>Dnmt 1</i> (two copies) (Robinson et al., 2011)	3.1
	Isoform 1, partial: CCA62565.1	
	Isoform 2, partial: CCA62566.1	
	<i>Dnmt 2</i> (Robinson et al., 2011)	3.2
	Partial sequence: CCA62567.1	
4. Methyl-CpG-binding domain (MBD):	<i>MBD-like</i>	4.1
5. Histone methyltransferases (HMT):	<i>Set</i>	5.1
	<i>MLL</i>	5.2
	<i>Enhancer of zeste</i>	5.3
	<i>Buzidau</i>	5.4
	<i>Smyd4</i>	5.5
	<i>G9a</i> : not found	
	<i>K9H9</i> : not found	
6. Histone demethylases (HDM):	Su(var)3-3	6.1
	Su(var)3-9	6.2